arXiv:math/0411333v2 [math.PR] 25 Feb 2005

THE EMPIRICAL DISTRIBUTION OF THE EIGENVALUES OF A GRAM MATRIX WITH A GIVEN VARIANCE PROFILE

W. HACHEM, P. LOUBATON AND J. NAJIM

Abstract. Consider a $N \times n$ random matrix $Y_n = (Y_{ij}^n)$ where the entries are given by $Y_{ij}^n = \frac{\sigma(i/N,j/n)}{\sqrt{n}} X_{ij}^n$, the X_{ij}^n being centered i.i.d. and $\sigma : [0,1]^2 \to (0,\infty)$ being a continuous function called a variance profile. Consider now a deterministic $N \times n$ matrix $\Lambda_n = \left(\Lambda_{ij}^n\right)$ whose non diagonal elements are zero. Denote by Σ_n the non-centered matrix $Y_n + \Lambda_n$ and by $N \wedge n = \min(N,n)$. Then under the assumption that $\lim_{n\to\infty} \frac{N}{n} = c > 0$ and

$$\frac{1}{N \wedge n} \sum_{i=1}^{N\wedge n} \delta_{\left(\frac{i}{N\wedge n}, (\Lambda_{ii}^n)^2\right)} \xrightarrow[n\to\infty]{} H(dx, d\lambda),$$

where H is a probability measure, it is proven that the empirical distribution of the eigenvalues of $\Sigma_n \Sigma_n^T$ converges almost surely in distribution to a non random probability measure. This measure is characterized in terms of its Stieltjes transform, which is obtained with the help of an auxiliary system of equations. This kind of results is of interest in the field of wireless communication.

Résumé. Soit $Y_n = (Y_{ij}^n)$ une matrice $N \times n$ dont les entrées sont données par $Y_{ij}^n = \frac{\sigma(i/N,j/n)}{\sqrt{n}} X_{ij}^n$, les X_{ij}^n étant des variables aléatoires centrées, i.i.d. et où $\sigma : [0,1]^2 \to (0,\infty)$ est une fonction continue qu'on appellera profil de variance. Considérons une matrice déterministe $\Lambda_n = \left(\Lambda_{ij}^n\right)$ de dimensions $N \times n$ dont les éléments non diagonaux sont nuls. Appelons Σ_n la matrice non centrée définie par $\Sigma_n = Y_n + \Lambda_n$ et notons $N \wedge n = \min(N,n)$. Sous les hypothèses que $\lim_{n\to\infty} \frac{N}{n} = c > 0$ et que

$$\frac{1}{N \wedge n} \sum_{i=1}^{N\wedge n} \delta_{\left(\frac{i}{N\wedge n}, (\Lambda_{ii}^n)^2\right)} \xrightarrow[n\to\infty]{} H(dx, d\lambda),$$

où H est une probabilité, on démontre que la mesure empirique des valeurs propres de $\Sigma_n \Sigma_n^T$ converge presque sûrement vers une mesure de probabilité déterministe. Cette mesure est caractérisée par sa transformée de Stieltjes, qui s'obtient à l'aide d'un système d'équations auxiliaire. Ce type de résultats présente un intérêt dans le champ des communications numériques sans fil.

Key words and phrases: Random Matrix, empirical distribution of the eigenvalues, Stieltjes transform.
AMS 2000 subject classification: Primary 15A52, Secondary 15A18, 60F15.

1. Introduction

Consider a $N \times n$ random matrix $Y_n = (Y_{ij}^n)$ where the entries are given by

$$Y_{ij}^n = \frac{\sigma(i/N, j/n)}{\sqrt{n}} X_{ij}^n \tag{1.1}$$

Date: February 1, 2008.

where $\sigma : [0,1] \times [0,1] \to (0,\infty)$ is a continuous function called a variance profile and the random variables X_{ij}^n are real, centered, independent and identically distributed (i.i.d.) with finite $4+\epsilon$ moment. Consider a real deterministic $N \times n$ matrix $\Lambda_n = (\Lambda_{ij}^n)$ whose non-diagonal elements are zero and consider the matrix $\Sigma_n = Y_n + \Lambda_n$. This model has two interesting features: The random variables are independent but not i.i.d. since the variance may vary and Λ_n, the centering perturbation of Y_n, though (pseudo) diagonal can be of full rank. The purpose of this article is to study the convergence of the empirical distribution of the eigenvalues of the Gram random matrix $\Sigma_n \Sigma_n^T$ (Σ_n^T being the transpose of Σ_n) when $n \to +\infty$ and $N \to +\infty$ in such a way that $\frac{N}{n} \to c$, $0 < c < +\infty$.

The asymptotics of the spectrum of $N \times N$ Gram random matrices $Z_n Z_n^T$ have been widely studied in the case where Z_n is centered (see Marčenko and Pastur [14], Yin [21], Silverstein et al. [15, 16], Girko [8, 9], Khorunzhy et al. [12], Boutet de Monvel et al. [4], etc.). For an overview on asymptotic spectral properties of random matrices, see Bai [2]. The case of a Gram matrix $Z_n Z_n^T$ where Z_n is non centered has comparatively received less attention. Let us mention Girko ([9], chapter 7) where a general study is carried out for the matrix $Z_n = (W_n + A_n)$ where W_n has a given variance profile and A_n is deterministic. In [9], it is proved that the entries of the resolvent $(Z_n Z_n^T - zI)^{-1}$ have the same asymptotic behavior as the entries of a certain deterministic holomorphic $N \times N$ matrix valued function $T_n(z)$. This matrix-valued function is characterized by a non linear system of $(n+N)$ coupled functional equations. Using different methods, Brent Dozier and Silverstein [5] study the eigenvalue asymptotics of the matrix $(R_n+X_n)(R_n+X_n)^T$ in the case where the matrices X_n and R_n are independent random matrices, X_n has i.i.d. entries and the empirical distribution of $R_n R_n^T$ converges to a non-random distribution. It is proved there that the eigenvalue distribution of $(R_n+X_n)(R_n+X_n)^T$ converges almost surely towards a deterministic distribution whose Stieltjes transform is uniquely defined by a certain functional equation.

As in [5], the model studied in this article, i.e. $\Sigma_n = Y_n + \Lambda_n$, is a particular case of the general case studied in ([9], chapter 7, equation K_7) for which there exists a limiting distribution for the empirical distribution of the eigenvalues. Since the centering term Λ_n is pseudo-diagonal, the proof of the convergence of the empirical distribution of the eigenvalues is based on a direct analysis of the diagonal terms of the resolvent $(\Sigma_n \Sigma_n^T - zI)^{-1}$. This analysis leads in a natural way to the equations characterizing the Stieltjes transform of the limiting probability distribution of the eigenvalues. In the Wigner case with a variance profile (matrix Y_n and the variance profile are symmetric), let us mention the recent work of Anderson and Zeitouni [1] where the asymptotics of the spectum and central limit theorems are investigated by means of systematic combinatorial enumeration.

Recently, many of these results have been applied to the field of Signal Processing and Communication Systems and some new ones have been developed for that purpose (Silverstein and Combettes [17], Tse et al. [18, 19], Debbah et al. [7], Li et al. [13], etc.). The issue addressed in this paper is mainly motivated by the performance analysis of multiple-input multiple-output (MIMO) digital communication systems. In MIMO systems with n transmit antennas and N receive antennas, one can model the communication channel by a $N \times n$ matrix $H_n = (H_{ij}^n)$ where the entries H_{ij}^n represent the complex gain between transmit antenna i and receive antenna j. The statistics $C_n = \frac{1}{n} \log\det(I_n + \frac{H_n H_n^\star}{\sigma^2})$ (where $H_n^\star$ is the hermitian adjoint and σ^2 represents the variance of an additive noise corrupting the received signals) is a popular performance analysis index since it has been shown in information theory that C_n is the maximum number of bits per channel use and per antenna that can be transmitted reliably in a MIMO system with channel matrix H_n. Since

$$C_n = \frac{1}{n}\sum_{k=1}^{N} \log(1+\frac{\mu_k}{\sigma^2}),$$

where $(\mu_k)_{1\le k\le N}$ are the eigenvalues of $H_n H_n^\star$, the empirical distribution of the eigenvalues of $H_n H_n^\star$ gives direct information on C_n (see Tulino and Verdu [20] for an exhaustive review of recent results). For wireless systems, matrix H_n is often modelled as a zero-mean Gaussian random matrix and several articles have recently been devoted to the study of the impact of the channel statistics (via the eigenvalues of $H_n H_n^\star$) on the probability distribution of C_n (Chuah et al. [6], Goldsmith et al. [10], see also [20] and the references therein). Of particular interest is also the channel matrix $H_n = F_N(Y_n + \Lambda_n)F_n^T$ where $F_k = (F_{pq}^k)_{1\le p,q\le k}$ is the Fourier matrix (i.e. $F_{pq}^k = k^{-\frac{1}{2}}\exp(2i\pi\frac{(p-1)(q-1)}{k})$) and the matrix Y_n is given by (1.1) (see [20], p. 139 for more details). The matrices H_n and Σ_n having the same singular values, we will focus on the study of the empirical distribution of the singular values of Σ_n. Moreover, we will focus on matrices with real entries since the complex case is a straightforward extension.

In the sequel, we will study simultaneously quantities (Stieltjes kernels) related to the Stieltjes transforms of $\Sigma_n\Sigma_n^T$ and $\Sigma_n^T\Sigma_n$. Even if the Stieltjes transforms of $\Sigma_n\Sigma_n^T$ and $\Sigma_n^T\Sigma_n$ are related in an obvious way, the corresponding Stieltjes kernels are not, as we shall see. We will prove that if $N/n \xrightarrow[n\to\infty]{} c > 0$ (since we study at the same time $\Sigma_n\Sigma_n^T$ and $\Sigma_n^T\Sigma_n$, we assume without loss of generality that $c \le 1$) and if there exists a probability measure H on $[0,1]\times\mathbb{R}$ with compact support such that

$$\frac{1}{N}\sum_{i=1}^{N}\delta_{\left(\frac{i}{N},(\Lambda_{ii}^n)^2\right)} \xrightarrow[n\to\infty]{\mathcal{D}} H(dx,d\lambda)$$

where $\mathcal{D}$ stands for the convergence in distribution, then almost surely, the empirical distribution of the eigenvalues of the random matrix $\Sigma_n\Sigma_n^T$ (resp. $\Sigma_n^T\Sigma_n$) converges in distribution to a deterministic probability distribution $\mathbb{P}$ (resp. $\tilde{\mathbb{P}}$). The probability distributions $\mathbb{P}$ and $\tilde{\mathbb{P}}$ are characterized in terms of their Stieltjes transform

$$f(z) = \int_{\mathbb{R}^+}\frac{\mathbb{P}(dx)}{x-z} \quad \text{and} \quad \tilde{f}(z) = \int_{\mathbb{R}^+}\frac{\tilde{\mathbb{P}}(dx)}{x-z}, \quad \mathrm{Im}(z)\neq 0.$$

as follows. Consider the following system of equations

$$\int g\,d\pi_z = \int \frac{g(u,\lambda)}{-z(1+\int\sigma^2(u,\cdot)d\tilde{\pi}_z) + \frac{\lambda}{1+c\int\sigma^2(\cdot,cu)d\pi_z}} H(du,d\lambda)$$

$$\begin{aligned}\int g\,d\tilde{\pi}_z = c\int &\frac{g(cu,\lambda)}{-z(1+c\int\sigma^2(\cdot,cu)d\pi_z) + \frac{\lambda}{1+\int\sigma^2(u,\cdot)d\tilde{\pi}_z}} H(du,d\lambda)\\ &+ (1-c)\int_c^1 \frac{g(u,0)}{-z(1+c\int\sigma^2(\cdot,u)d\pi_z)}\,du\end{aligned}$$

where the unknown parameters are the complex measures π_z and $\tilde{\pi}_z$. Both equalities stand for every continuous function $g:\mathcal{H}\to\mathbb{R}$, where $\mathcal{H}\subset[0,1]\times\mathbb{R}$ is the compact support of H. Then, this system admits a unique pair of solutions $(\pi_z(dx,d\lambda),\tilde{\pi}_z(dx,d\lambda))$. In particular,

π_z is absolutely continuous with respect to H while $\tilde{\pi}_z$ is not (see Remarks 2.5, 2.6 and 2.7 for more details). The Stieltjes transforms f and $\tilde{f}$ are then given by

$$f(z) = \int_{[0,1]\times\mathbb{R}} \pi_z(dx, d\lambda) \quad \text{and} \quad \tilde{f}(z) = \int_{[0,1]\times\mathbb{R}} \tilde{\pi}_z(dx, d\lambda).$$

The article is organized as follows. In Section 2, the notations and the assumptions are introduced and the main results (Theorem 2.3 and Theorem 2.4) are stated. Section 3 is devoted to the proof of Theorem 2.3. Section 4 is devoted to the proof of Theorem 2.4. In Section 5, we state as corollaries of Theorems 2.3 and 2.4 two simpler results which already exist in the literature. We also give details for the statement of the results in the complex setting (Section 5.3).

2. Convergence of the Stieltjes Transform

2.1. Notations and Assumptions.

Let $N = N(n)$ be a sequence of integers such that

$$\lim_{n\to\infty} \frac{N(n)}{n} = c.$$

Consider a $N \times n$ random matrix Y_n where the entries are given by

$$Y_{ij}^n = \frac{\sigma(i/N, j/n)}{\sqrt{n}} X_{ij}^n$$

where X_{ij}^n and σ are defined below.

Assumption A-1. *The random variables* $(X_{ij}^n \ ; \ 1 \le i \le N, 1 \le j \le n \, , n \ge 1)$ *are real, independent and identically distributed. They are centered with* $\mathbb{E}(X_{ij}^n)^2 = 1$ *and satisfy:*

$$\exists\, \epsilon > 0, \quad \mathbb{E}|X_{ij}^n|^{4+\epsilon} < \infty.$$

where $\mathbb{E}$ denotes the expectation.

Assumption A-2. *The real function* $\sigma : [0,1] \times [0,1] \to \mathbb{R}$ *is such that* σ^2 *is continuous. Therefore there exist a non-negative constant* $\sigma_{\max}$ *such that*

$$\forall (x,y) \in [0,1]^2, \quad 0 \le \sigma^2(x,y) \le \sigma_{\max}^2 < \infty. \tag{2.1}$$

Remark 2.1. As long as σ^2 remains continuous, σ can vanish on portions of the domain $[0,1]^2$.

Denote by $\mathrm{var}(Z)$ the variance of the random variable Z. Since $\mathrm{var}(Y_{ij}^n) = \sigma^2(i/N, j/n)/n$, the function σ will be called a variance profile. Denote by $\delta_{z_0}(\,dz)$ the dirac measure at point z_0. The indicator function of A is denoted by $1_A(x)$. Denote by $C_b(\mathcal{X})$ (resp. $C_b(\mathcal{X};\mathbb{C})$) the set of real (resp. complex) continuous and bounded functions over the topological set $\mathcal{X}$ and by $\|f\|_\infty = \sup_{x\in\mathcal{X}} |f(x)|$, the supremum norm. If $\mathcal{X}$ is compact, we simply write $C(\mathcal{X})$ (resp. $C(\mathcal{X};\mathbb{C})$)) instead of $C_b(\mathcal{X})$ (resp. $C_b(\mathcal{X};\mathbb{C})$)).

We will denote by $\xrightarrow{\mathcal{D}}$ the convergence in distribution for probability measures and by $\xrightarrow{w}$ the weak convergence for bounded complex measures.

Consider a real deterministic $N \times n$ matrix $\Lambda_n = (\Lambda_{ij}^n)$ whose non-diagonal entries are zero. We will often write Λ_{ij} instead of Λ_{ij}^n. We introduce the $N \times n$ matrix $\Sigma_n = Y_n + \Lambda_n$. For every matrix A, we will denote by A^T its transpose, by $\mathrm{Tr}(A)$ its trace (if A is square) and by $F^{A\,A^T}$, the empirical distribution function of the eigenvalues of $A\,A^T$. Since we will study at the same time the limiting spectrum of the matrices $\Sigma_n\Sigma_n^T$ and $\Sigma_n^T\Sigma_n$, we can assume without loss of generality that $c \le 1$. We also assume for simplicity that $N \le n$.

We assume that:

Assumption A-3. *There exists a probability measure $H(\,du, d\lambda)$ over the set $[0,1]\times\mathbb{R}$ with compact support $\mathcal{H}$ such that*

$$\frac{1}{N}\sum_{i=1}^{N}\delta_{\left(\frac{i}{N},\,(\Lambda_{ii}^n)^2\right)}(du,d\lambda) \xrightarrow[n\to\infty]{\mathcal{D}} H(\,du,d\lambda). \tag{2.2}$$

Remark 2.2 (The complex case). Assumptions (A-1), (A-2) and (A-3) must be slightly modified in the complex setting. Related modifications are stated in Section 5.3.

When dealing with vectors, the norm $\|\cdot\|$ will denote the Euclidean norm. In the case of matrices, the norm $\|\cdot\|$ will refer to the spectral norm.

Remark 2.3. Due to (A-3), we can assume without loss of generality that the Λ_{ii}^n's are bounded for n large enough. In fact, suppose not, then by (A-3), $\frac{1}{N}\sum_{i=1}^N \delta_{\Lambda_{ii}^2} \to H_\Lambda(d\lambda)$ whose support is compact and, say, included in $[0,K]$. Then Portmanteau's theorem yields $\frac{1}{N}\sum_1^N 1_{[0,\ K+\delta]}(\Lambda_{ii}^2) \to 1$ thus

$$\frac{\#\{\Lambda_{ii},\ \Lambda_{ii}^2 \notin [0,\ K+\delta]\}}{N} = 1 - \frac{1}{N}\sum_{1}^{N} 1_{[0,\ K+\delta]}(\Lambda_{ii}^2) \xrightarrow[n\to\infty]{} 0. \tag{2.3}$$

Denote by $\check{\Lambda}_n = (\check{\Lambda}_{ij}^n)$ the matrix whose non-diagonal elements are zero and set $\check{\Lambda}_{ii}^n = \Lambda_{ii}^n 1_{\{(\Lambda_{ii}^n)^2 \le K+\delta\}}$. Then it is straightforward to check that $\frac{1}{N}\sum_{i=1}^N \delta_{\left(\frac{i}{N},\,\check{\Lambda}_{ii}^2\right)}(du,d\lambda) \to H(\,du,d\lambda)$. Moreover, if $\check{\Sigma}_n = Y_n + \check{\Lambda}_n$ then

$$\|F^{\Sigma\Sigma^T} - F^{\check{\Sigma}\check{\Sigma}^T}\|_\infty \overset{(a)}{\le} \frac{\mathrm{rank}(\check{\Sigma}-\Sigma)}{N} \overset{(b)}{\le} \frac{\#\{\Lambda_{ii},\ \Lambda_{ii}^2 \notin [\ 0,\ K+\delta]\}}{N} \xrightarrow[n\to\infty]{(c)} 0.$$

where (a) follows from Lemma 3.5 in [21] (see also [16], Section 2), (b) follows from the fact that for a rectangular matrix A, rank(A) $\le$ the number of non zero entries of A and (c) follows from (2.3). Therefore, $F^{\Sigma\Sigma^T}$ converges iff $F^{\check{\Sigma}\check{\Sigma}^T}$ converges. In this case they share the same limit. Remark 2.3 is proved.

Remark 2.4. Due to Remark 2.3, we will assume in the sequel that for all n, the support of $\frac{1}{N}\sum \delta_{\left(\frac{i}{N},\,\Lambda_{ii}^2\right)}$ is included in a compact set $\mathcal{K} \subset [0,1]\times\mathbb{R}$.

Let $\mathbb{C}^+ = \{z \in \mathbb{C},\ \mathrm{Im}(z) > 0\}$ and $\mathbb{C}^\nabla = \{z \in \mathbb{C}^+,\ |\mathrm{Re}(z)| \le \mathrm{Im}(z)\}$.

2.2. Stieltjes transforms and Stieltjes kernels.

Let ν be a bounded nonnegative measure over $\mathbb{R}$. Its Stieltjes transform f is defined by:

$$f(z) = \int_{\mathbb{R}} \frac{\nu(d\lambda)}{\lambda - z}, \quad z \in \mathbb{C}^+.$$

We list below the main properties of the Stieltjes transforms that will be needed in the sequel.

Proposition 2.1. *The following properties hold true:*

1. *Let f be the Stieltjes transform of ν, then*
   - *the function f is analytic over $\mathbb{C}^+$,*
   - *the function f satisfies: $|f(z)| \leq \frac{\nu(\mathbb{R})}{\mathrm{Im}(z)}$,*
   - *if $z \in \mathbb{C}^+$ then $f(z) \in \mathbb{C}^+$,*
   - *if $\nu(-\infty, 0) = 0$ then $z \in \mathbb{C}^+$ implies $z\, f(z) \in \mathbb{C}^+$.*

2. *Conversely, let f be a function analytic over $\mathbb{C}^+$ such that $f(z) \in \mathbb{C}^+$ if $z \in \mathbb{C}^+$ and $|f(z)||Im(z)|$ bounded on $\mathbb{C}^+$. Then, f is the Stieltjes transform of a bounded positive measure μ and $\mu(\mathbb{R})$ is given by*
   $$\mu(\mathbb{R}) = \lim_{y \to +\infty} -iy\, f(iy).$$
   If moreover $zf(z) \in \mathbb{C}^+$ if $z \in \mathbb{C}^+$ then, $\mu(\mathbb{R}^-) = 0$.

3. *Let $\mathbb{P}_n$ and $\mathbb{P}$ be probability measures over $\mathbb{R}$ and denote by f_n and f their Stieltjes transforms. Then*
   $$\left(\forall z \in \mathbb{C}^+,\ f_n(z) \xrightarrow[n \to \infty]{} f(z) \right) \quad \Rightarrow \quad \mathbb{P}_n \xrightarrow[n \to \infty]{\mathcal{D}} \mathbb{P}.$$

Let A be an $n \times p$ matrix and let I_n be the $n \times n$ identity. The resolvent of AA^T is defined by

$$Q(z) = (AA^T - z\, I_n)^{-1} = (\rho_{ij}(z))_{1 \leq i,j, \leq n}\,, \quad z \in \mathbb{C} \backslash \mathbb{R}.$$

The following properties are straightforward.

Proposition 2.2. *Let $Q(z)$ be the resolvent of AA^T, then:*

1. *For all $z \in \mathbb{C}^+$, $\|Q(z)\| \leq (\mathrm{Im}(z))^{-1}$. Similarly, $|\rho_{ij}(z)| \leq (\mathrm{Im}(z))^{-1}$.*

2. *The function $h_n(z) = \frac{1}{n} \mathrm{Tr}\ Q(z)$ is the Stieltjes transform of the empirical distribution probability associated to the eigenvalues of AA^T. Since these eigenvalues are nonnegative, $z\, h_n(z) \in \mathbb{C}^+$ for $z \in \mathbb{C}^+$.*

3. *Let $\vec{\xi}$ be a $n \times 1$ vector, then $\mathrm{Im} \left(z \vec{\xi} Q(z) \vec{\xi}^T \right) \in \mathbb{C}^+$ for $z \in \mathbb{C}^+$.*

Denote by $\mathcal{M}_{\mathbb{C}}(\mathcal{X})$ the set of complex measures over the topological set $\mathcal{X}$. In the sequel, we will call Stieltjes kernel every application

$$\mu : \mathbb{C}^+ \to \mathcal{M}_{\mathbb{C}}(\mathcal{X})$$

either denoted $\mu(z, dx)$ or $\mu_z(dx)$ and satisfying:

1. $\forall g \in C_b(\mathcal{X}),\ \int g\, d\mu_z$ is analytic over $\mathbb{C}^+$,
2. $\forall z \in \mathbb{C}^+,\ \forall g \in C_b(\mathcal{X})$,
   $$\left| \int g\, d\mu_z \right| \leq \frac{\|g\|_\infty}{\mathrm{Im}(z)}$$
3. $\forall z \in \mathbb{C}^+,\ \forall g \in C_b(\mathcal{X})$ and $g \geq 0$ then $\mathrm{Im} \left(\int g\, d\mu_z \right) \geq 0$,
4. $\forall z \in \mathbb{C}^+,\ \forall g \in C_b(\mathcal{X})$ and $g \geq 0$ then $\mathrm{Im} \left(z \int g\, d\mu_z \right) \geq 0$.

Let us introduce the following resolvents:

$$\begin{aligned} Q_n(z) &= (\Sigma_n \Sigma_n^T - zI_N)^{-1} = (q_{ij}(z))_{1\le i,j\le N}\,, \quad z \in \mathbb{C}^+, \\ \tilde{Q}_n(z) &= (\Sigma_n^T \Sigma_n - zI_n)^{-1} = (\tilde{q}_{ij}(z))_{1\le i,j\le n}\,, \quad z \in \mathbb{C}^+, \end{aligned}$$

and the following empirical measures defined for $z \in \mathbb{C}^+$

$$L_z^n(du, d\lambda) = \frac{1}{N}\sum_{i=1}^N q_{ii}(z)\ \delta_{\left(\frac{i}{N},\Lambda_{ii}^2\right)}(du, d\lambda), \tag{2.4}$$

$$\begin{aligned} \tilde{L}_z^n(du, d\lambda) &= \frac{1}{n}\sum_{i=1}^N \tilde{q}_{ii}(z)\ \delta_{\left(\frac{i}{n},\Lambda_{ii}^2\right)}(du, d\lambda) \\ &\quad + \left(\frac{1}{n}\sum_{i=N+1}^n \tilde{q}_{ii}(z)\ \delta_{\frac{i}{n}}(du)\otimes\delta_0(d\lambda)\right) 1_{\{N<n\}}, \end{aligned} \tag{2.5}$$

where $\otimes$ denotes the product of measures. Since $q_{ii}(z)$ (resp. $\tilde{q}_{ii}(z)$) is analytic over $\mathbb{C}^+$, satisfies $|q_{ii}(z)| \le (\mathrm{Im}(z))^{-1}$ and $\min\left(\mathrm{Im}(q_{ii}(z)), \mathrm{Im}(zq_{ii}(z))\right) > 0$, L^n (resp. $\tilde{L}^n$) is a Stieltjes kernel. Recall that due to Remark 2.4, L^n and $\tilde{L}^n$ have supports included in the compact set $\mathcal{K}$.

Remark 2.5 (on the limiting support of L^n)*.* Consider a converging subsequence of L_z^n, then its limiting support is necessarily included in $\mathcal{H}$.

Remark 2.6 (on the limiting support of $\tilde{L}^n$)*.* Denote by H_c the image of the probability measure H under the application $(u,\lambda)\mapsto(cu,\lambda)$, by $\mathcal{H}_c$ its support, by $\mathcal{R}$ the support of the measure $1_{[c,1]}(u)\,du\otimes\delta_0(d\lambda)$. Let $\tilde{\mathcal{H}} = \mathcal{H}_c \cup \mathcal{R}$. Notice that $\tilde{\mathcal{H}}$ is obviously compact. Consider a converging subsequence of $\tilde{L}_z^n$, then its limiting support is necessarily included in $\tilde{\mathcal{H}}$.

2.3. Convergence of the empirical measures L_z^n and $\tilde{L}_z^n$.

Theorem 2.3. *Assume that H is a probability measure over the set $[0,1]\times\mathbb{R}$ with compact support $\mathcal{H}$. Assume moreover that (A-2) holds. Then the system of equations*

$$\int g\,d\pi_z = \int \frac{g(u,\lambda)}{-z(1+\int\sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta)) + \frac{\lambda}{1+c\int\sigma^2(t,cu)\pi(z,dt,d\zeta)}} H(du,d\lambda) \tag{2.6}$$

$$\begin{aligned} \int g\,d\tilde{\pi}_z &= c\int \frac{g(cu,\lambda)}{-z(1+c\int\sigma^2(t,cu)\pi(z,dt,d\zeta)) + \frac{\lambda}{1+\int\sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta)}} H(du,d\lambda) \\ &\quad + (1-c)\int_c^1 \frac{g(u,0)}{-z(1+c\int\sigma^2(t,u)\pi(z,dt,d\zeta))}\,du \end{aligned} \tag{2.7}$$

where (2.6) and (2.7) hold for every $g\in C(\mathcal{H})$, admits a unique couple of solutions $(\pi(z,dt,d\lambda),\tilde{\pi}(z,dt,d\lambda))$ among the set of Stieltjes kernels for which the support of measure π_z is included in $\mathcal{H}$ and the support of measure $\tilde{\pi}_z$ is included in $\tilde{\mathcal{H}}$.

Moreover the functions $f(z) = \int d\pi_z$ and $\tilde{f}(z) = \int d\tilde{\pi}_z$ are the Stieltjes transforms of probability measures.

Remark 2.7 (on the absolute continuity of π_z and $\tilde{\pi}_z$). Due to (2.6), the complex measure π_z is absolutely continuous with respect to H. However, it is clear from (2.7) that $\tilde{\pi}_z$ has an absolutely continuous part with respect to H_c (recall that H_c is the image of H under $(u,\lambda) \mapsto (cu,\lambda)$) and an absolutely continuous part with respect to $1_{[c,1]}(u)\,du \otimes \delta_0(d\lambda)$ (which is in general singular with respect to H_c). Therefore, it is much more convenient to work with Stieltjes kernels π and $\tilde{\pi}$ rather than with measure densities indexed by z.

Theorem 2.4. *Assume that (A-1), (A-2) and (A-3) hold and denote by π and $\tilde{\pi}$ the two Stieltjes kernels solutions of the coupled equations (2.6) and (2.7). Then*

(1) *Almost surely, the Stieltjes kernel L^n defined by (2.4) converges weakly to π, that is:*

$$a.s. \quad \forall z \in \mathbb{C}^+, \quad L_z^n \xrightarrow[n\to\infty]{w} \pi_z.$$

(2) *Almost surely, the Stieltjes kernel $\tilde{L}^n$ defined by (2.5) converges weakly to $\tilde{\pi}$.*

Proofs of Theorems 2.3 and 2.4 are postponed to Sections 3 and 4.

Corollary 2.5. *Assume that (A-1), (A-2) and (A-3) hold and denote by π and $\tilde{\pi}$ the two Stieltjes kernels solutions of the coupled equations (2.6) and (2.7). Then the empirical distribution of the eigenvalues of the matrix $\Sigma_n\Sigma_n^T$ converges almost surely to a non-random probability measure $\mathbb{P}$ whose Stieltjes transform $f(z) = \int_{\mathbb{R}^+} \frac{\mathbb{P}(dx)}{x-z}$ is given by:*

$$f(z) = \int_{\mathcal{H}} \pi_z(dx, d\lambda).$$

Similarly, the empirical distribution of the eigenvalues of the matrix $\Sigma_n^T\Sigma_n$ converges almost surely to a non-random probability measure $\tilde{\mathbb{P}}$ whose Stieltjes transform $\tilde{f}(z)$ is given by:

$$\tilde{f}(z) = \int_{\tilde{\mathcal{H}}} \tilde{\pi}_z(dx, d\lambda).$$

Proof of Corollary 2.5. The Stieltjes transform of $\Sigma_n\Sigma_n^T$ is equal to $\frac{1}{N}\sum_{i=1}^N q_{ii}(z) = \int dL_z^n$. By Theorem 2.4,

$$\text{a.s.} \quad \forall z \in \mathbb{C}^+, \quad \int dL_z^n \xrightarrow[n\to\infty]{} \int d\pi_z. \tag{2.8}$$

Since $\int d\pi_z$ is the Stieltjes transform of a probability measure $\mathbb{P}$ by Theorem 2.3, eq. (2.8) implies that $F^{\Sigma_n\Sigma_n^T}$ converges weakly to $\mathbb{P}$. One can similarly prove that $F^{\Sigma_n^T\Sigma_n}$ converges weakly to a probability measure $\tilde{\mathbb{P}}$. □

3. Proof of Theorem 2.3

We first introduce some notations. Denote by

$$\begin{aligned}
D(\tilde{\pi}_z,\pi_z)(u,\lambda) &= -z\left(1+\int \sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta)\right)+\frac{\lambda}{1+c\int \sigma^2(t,cu)\pi(z,dt,d\zeta)},\\
d(\pi_z)(u) &= 1+c\int \sigma^2(t,cu)\pi(z,dt,d\zeta),\\
\tilde{D}(\pi_z,\tilde{\pi}_z)(u,\lambda) &= -z\left(1+c\int \sigma^2(t,cu)\pi(z,dt,d\zeta)\right)+\frac{\lambda}{1+\int \sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta)},\\
\tilde{d}(\tilde{\pi}_z)(u) &= 1+\int \sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta),\\
\kappa(\pi_z)(u) &= -z\left(1+c\int \sigma^2(t,u)\pi(z,dt,d\zeta)\right).
\end{aligned}$$

Let ν be a complex measure over the set $\mathcal{H}$ (recall that $\mathcal{H}$ is compact by (A-3)) then we denote by $\|\nu\|_{\mathrm{tv}}$ the total variation norm of ν, that is

$$\begin{aligned}
\|\nu\|_{\mathrm{tv}} &= |\nu|(\mathcal{H})\\
&= \sup\left\{\left|\int f\,d\nu\right|,\ f\in C(\mathcal{H};\mathbb{C}),\ \|f\|_\infty\le 1\right\}.
\end{aligned}$$

3.1. **Proof of the unicity of the solutions.** Notice that the system of equations (2.6) and (2.7) remains true for every $g\in C(\mathcal{H};\mathbb{C})$ (consider $g=h+ik$) and assume that both $(\pi_z,\tilde{\pi}_z)$ and $(\rho_z,\tilde{\rho}_z)$ are pairs of solutions of the given system. Let $g\in C(\mathcal{H};\mathbb{C})$, then (2.6) yields:

$$\begin{aligned}
\int g\,d\pi_z-\int g\,d\rho_z &= \int \frac{z\,g(u,\lambda)\int\sigma^2(u,t)(\tilde{\pi}(z,dt,d\zeta)-\tilde{\rho}(z,dt,d\zeta))}{D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)}H(du,d\lambda)\\
&\quad+\int \frac{c\lambda\,g(u,\lambda)\int\sigma^2(t,cu)(\rho(z,dt,d\zeta)-\pi(z,dt,d\zeta))}{D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)\times d(\pi_z)\times d(\rho_z)}H(du,d\lambda)
\end{aligned}$$

and

$$\begin{aligned}
\left|\int g\,d\pi_z-\int g\,d\rho_z\right| &\le |z|\sigma^2_{\max}\,\|g\|_\infty\,\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}}\int\frac{dH}{|D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)|}\\
&\quad+c\,\sigma^2_{\max}\,\|g\|_\infty\,\|\pi_z-\rho_z\|_{\mathrm{tv}}\int\frac{\lambda H(du,d\lambda)}{|D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)\times d(\pi_z)\times d(\rho_z)|}
\end{aligned}$$

If one takes the supremum over the functions $g\in C(\mathcal{H};\mathbb{C})$, $\|g\|_\infty\le 1$, one gets :

$$\|\pi_z-\rho_z\|_{\mathrm{tv}}\le\boldsymbol{\alpha}\|\pi_z-\rho_z\|_{\mathrm{tv}}+\boldsymbol{\beta}\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}}$$

where

$$\begin{aligned}
\boldsymbol{\alpha}=\boldsymbol{\alpha}(\pi,\rho,\tilde{\pi},\tilde{\rho}) &= c\,\sigma^2_{\max}\int\frac{\lambda H(du,d\lambda)}{|D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)\times d(\pi_z)\times d(\rho_z)|}\\
\boldsymbol{\beta}=\boldsymbol{\beta}(\pi,\rho,\tilde{\pi},\tilde{\rho}) &= |z|\sigma^2_{\max}\int\frac{dH}{|D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)|}
\end{aligned}$$

Similarly, (2.7) yields:

$$\begin{aligned}
\int g\,d\tilde{\pi}_z - \int g\,d\tilde{\rho}_z &= c\int \frac{c\,z\,g(u,\lambda)\int \sigma^2(t,cu)(\pi(z,dt,d\zeta)-\rho(z,dt,d\zeta))}{\tilde{D}(\pi_z,\tilde{\pi}_z)\times\tilde{D}(\rho_z,\tilde{\rho}_z)} H(du,d\lambda) \\
&\quad + c\int \frac{\lambda\,g(u,\lambda)\int \sigma^2(u,t)(\tilde{\pi}(z,dt,d\zeta)-\tilde{\rho}(z,dt,d\zeta))}{\tilde{D}(\pi_z,\tilde{\pi}_z)\times\tilde{D}(\rho_z,\tilde{\rho}_z)\times\tilde{d}(\tilde{\pi}_z)\times\tilde{d}(\tilde{\rho}_z)} H(du,d\lambda) \\
&\quad + (1-c)\int_c^1 \frac{c\,z\,g(u,0)\int \sigma^2(t,u)(\pi(z,dt,d\zeta)-\rho(z,dt,d\zeta))}{\kappa(\pi_z)\times\kappa(\rho_z)} du
\end{aligned}$$

and

$$\begin{aligned}
\left|\int g\,d\tilde{\pi}_z - \int g\,d\tilde{\rho}_z\right| &\le c^2\sigma^2_{\max}|z|\,\|g\|_\infty\|\pi_z-\rho_z\|_{\mathrm{tv}}\int \frac{dH}{\left|\tilde{D}(\pi_z,\tilde{\pi}_z)\times\tilde{D}(\rho_z,\tilde{\rho}_z)\right|} \\
&\quad + c\,\sigma^2_{\max}\|g\|_\infty\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}}\int \frac{\lambda H(du,d\lambda)}{\left|\tilde{D}(\pi_z,\tilde{\pi}_z)\times\tilde{D}(\rho_z,\tilde{\rho}_z)\times\tilde{d}(\tilde{\pi}_z)\times\tilde{d}(\tilde{\rho}_z)\right|} \\
&\quad + (1-c)c\sigma^2_{\max}|z|\,\|g\|_\infty\|\pi_z-\rho_z\|_{\mathrm{tv}}\int_c^1 \frac{du}{|\kappa(\pi_z)\times\kappa(\rho_z)|}
\end{aligned}$$

As previously, by taking the supremum over $g \in C(\mathcal{H};\mathbb{C})$, $\|g\|_\infty \le 1$, we get:

$$\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}} \le \tilde{\boldsymbol{\alpha}}\|\pi_z-\rho_z\|_{\mathrm{tv}} + \tilde{\boldsymbol{\beta}}\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}}$$

where

$$\begin{aligned}
\tilde{\boldsymbol{\alpha}} = \tilde{\boldsymbol{\alpha}}(\pi,\rho,\tilde{\pi},\tilde{\rho}) &= c\,\sigma^2_{\max}|z|\left(c\int \frac{dH}{\left|\tilde{D}(\pi_z,\tilde{\pi}_z)\times\tilde{D}(\rho_z,\tilde{\rho}_z)\right|} + (1-c)\int_c^1 \frac{du}{|\kappa(\pi_z)\times\kappa(\rho_z)|}\right) \\
\tilde{\boldsymbol{\beta}} = \tilde{\boldsymbol{\beta}}(\pi,\rho,\tilde{\pi},\tilde{\rho}) &= c\,\sigma^2_{\max}\int \frac{\lambda H(du,d\lambda)}{\left|\tilde{D}(\pi_z,\tilde{\pi}_z)\times\tilde{D}(\rho_z,\tilde{\rho}_z)\times\tilde{d}(\tilde{\pi}_z)\times\tilde{d}(\tilde{\rho}_z)\right|}
\end{aligned}$$

We end up with the following inequations:

$$\left\{\begin{array}{ll}
\|\pi_z-\rho_z\|_{\mathrm{tv}} & \le \boldsymbol{\alpha}\|\pi_z-\rho_z\|_{\mathrm{tv}} + \boldsymbol{\beta}\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}} \\
\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}} & \le \tilde{\boldsymbol{\alpha}}\|\pi_z-\rho_z\|_{\mathrm{tv}} + \tilde{\boldsymbol{\beta}}\|\tilde{\pi}_z-\tilde{\rho}_z\|_{\mathrm{tv}}
\end{array}\right. \tag{3.1}$$

Let us prove now that for $z \in \mathbb{C}^\nabla$ with $\mathrm{Im}(z)$ large enough, then $\boldsymbol{\alpha} < \frac{1}{2}$.

Since π and $\tilde{\pi}$ are assumed to be Stieltjes kernels, $\mathrm{Im}\left(z\int\sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta)\right) \ge 0$ and $\mathrm{Im}\left(\int\sigma^2(t,cu)\pi(z,dt,d\zeta)\right) \ge 0$. Therefore, $\mathrm{Im}(D(\tilde{\pi}_z,\pi_z)) \le -\mathrm{Im}(z)$ and hence $|\mathrm{Im}(D(\tilde{\pi}_z,\pi_z))| \ge \mathrm{Im}(z)$. Similarly, $|\mathrm{Im}(D(\tilde{\rho}_z,\rho_z))| \ge \mathrm{Im}(z)$. Thus,

$$\frac{1}{|D(\tilde{\pi}_z,\pi_z)\times D(\tilde{\rho}_z,\rho_z)|} \le \frac{1}{\mathrm{Im}^2(z)}.$$

Now consider $zd(\pi_z)$. As previously, $\mathrm{Im}(zd(\pi_z)) \ge \mathrm{Im}(z)$. As $|zd(\pi_z)| \ge |\mathrm{Im}(zd(\pi_z))|$, this implies that $\frac{1}{|zd(\pi_z)|} \le \frac{1}{\mathrm{Im}(z)}$ and $\frac{1}{|d(\pi_z)|} \le \frac{|z|}{\mathrm{Im}(z)}$. Since $z \in \mathbb{C}^\nabla$, $\frac{|z|}{\mathrm{Im}(z)} \le \sqrt{2}$. The same argument holds for $d(\rho_z)$ thus we get

$$\boldsymbol{\alpha} \le \frac{2\,c\,\sigma^2_{\max}\int\lambda H(du,d\lambda)}{\mathrm{Im}^2(z)} < \frac{1}{2} \quad \text{for } \mathrm{Im}(z) \text{ large enough.}$$

With similar arguments, one can prove that

$$\boldsymbol{\beta} \le \frac{\sqrt{2}\sigma_{\max}^2}{\mathrm{Im}(z)}, \quad \tilde{\boldsymbol{\alpha}} \le \frac{3\sigma_{\max}^2}{\mathrm{Im}(z)}, \quad \tilde{\boldsymbol{\beta}} \le \frac{2\sigma_{\max}^2}{\mathrm{Im}^2(z)} \int \lambda H(\,du d\lambda). \tag{3.2}$$

Therefore $\max(\boldsymbol{\alpha}, \boldsymbol{\beta}, \tilde{\boldsymbol{\alpha}}, \tilde{\boldsymbol{\beta}}) \le \theta < \frac{1}{2}$ for $z \in \mathbb{C}^\nabla$ and $\mathrm{Im}(z)$ large enough where θ does not depend on $(\pi, \tilde{\pi}, \rho, \tilde{\rho})$. Therefore, the system (3.1) yields

$$\|\pi_z - \rho_z\|_{\mathrm{tv}} = \|\tilde{\pi}_z - \tilde{\rho}_z\|_{\mathrm{tv}} = 0 \quad \text{for } z \in \mathbb{C}^\nabla \text{ and } \mathrm{Im}(z) \text{ large enough.}$$

Now take $z \in \mathbb{C}^+$ and $g \in C(\mathcal{H})$. Since $\int g\, d\pi_z$ and $\int g\, d\rho_z$ (resp. $\int g\, d\tilde{\pi}_z$ and $\int g\, d\tilde{\rho}_z$) are analytic over $\mathbb{C}^+$ and are equal in $\mathbb{C}^\nabla$ for $\mathrm{Im}(z)$ large enough, they are equal everywhere. Since this is true for all $g \in C(\mathcal{H})$, π_z and ρ_z (resp. $\tilde{\pi}_z$ and $\tilde{\rho}_z$) are identical on $\mathbb{C}^+$. This proves the unicity.

3.2. **Proof of the existence of solutions.** Let us now prove the existence of solutions to (3.1). Define by recursion

$$\pi^0(z, du, d\lambda) = \tilde{\pi}^0(z, du, d\lambda) = -\frac{1}{z} H(du, d\lambda),$$

and

$$\left\{ \begin{array}{ll} \int g(u,\lambda)\pi^p(z, du, d\lambda) & = \int \frac{g(u,\lambda)}{D(\tilde{\pi}_z^{p-1}, \pi_z^{p-1})} H(du, d\lambda) \\ \int g(u,\lambda)\tilde{\pi}^p(z, du, d\lambda) & = c \int \frac{g(cu,\lambda)}{\tilde{D}(\pi_z^{p-1}, \tilde{\pi}_z^{p-1})} H(du, d\lambda) + (1-c) \int_c^1 \frac{g(u,0)}{\kappa(\pi_z^{p-1})} du \end{array} \right.$$

for all $g \in C(\mathcal{H})$. It is straightforward to check that π_z^0 (resp. $\tilde{\pi}_z^0$) is a Stieltjes kernel. Moreover, this remains true for π^p and $\tilde{\pi}^p$ by induction over p. As for the unicity, we can establish

$$\left\{ \begin{array}{ll} \|\pi_z^p - \pi_z^{p-1}\|_{\mathrm{tv}} & \le \boldsymbol{\alpha} \|\pi_z^{p-1} - \pi_z^{p-2}\|_{\mathrm{tv}} + \boldsymbol{\beta} \|\tilde{\pi}_z^{p-1} - \tilde{\pi}_z^{p-2}\|_{\mathrm{tv}} \\ \|\tilde{\pi}_z^p - \tilde{\pi}_z^{p-1}\|_{\mathrm{tv}} & \le \tilde{\boldsymbol{\alpha}} \|\pi_z^{p-1} - \pi_z^{p-2}\|_{\mathrm{tv}} + \tilde{\boldsymbol{\beta}} \|\tilde{\pi}_z^{p-1} - \tilde{\pi}_z^{p-2}\|_{\mathrm{tv}} \end{array} \right. \tag{3.3}$$

where $\boldsymbol{\alpha}, \tilde{\boldsymbol{\alpha}}, \boldsymbol{\beta}$ and $\tilde{\boldsymbol{\beta}}$ depend on $(\pi_z^{p-1}, \tilde{\pi}_z^{p-1}, \pi_z^{p-2}, \tilde{\pi}_z^{p-2})$. As in (3.2), one can prove that

$$\max(\boldsymbol{\alpha}, \tilde{\boldsymbol{\alpha}}, \boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) \le \theta < \frac{1}{2} \quad \text{for } z \in \mathbb{C}^\nabla \text{ and } \mathrm{Im}(z) \text{ large enough}$$

where θ does not depend on $(\pi_z^{p-1}, \tilde{\pi}_z^{p-1}, \pi_z^{p-2}, \tilde{\pi}_z^{p-2})$. Therefore, (π_z^p) and $(\tilde{\pi}_z^p)$ are Cauchy sequences with respect to the norm $\|\cdot\|_{\mathrm{tv}}$ whenever $z \in \mathbb{C}^\nabla$ and $\mathrm{Im}(z)$ is large enough. This yields the existence and unicity of kernels π_z and $\tilde{\pi}_z$ such that

$$\forall g \in C(\mathcal{H}), \quad \left\{ \begin{array}{l} \int g\, d\pi_z^p \xrightarrow[p\to\infty]{} \int g\, d\pi_z \\ \int g\, d\tilde{\pi}_z^p \xrightarrow[p\to\infty]{} \int g\, d\tilde{\pi}_z \end{array} \right.$$

whenever $z \in \mathbb{C}^\nabla$ and $\mathrm{Im}(z)$ is large enough.

Let $g \in C(\mathcal{H})$ be fixed. Recall that $z \mapsto \int g\, d\pi_z^p$ is analytic on $\mathbb{C}^+$ and that $\forall z \in \mathbb{C}^+$, $|\int g\, d\pi_z^p| \le \frac{\|g\|_\infty}{\mathrm{Im}(z)}$. Therefore, $(\int g\, d\pi_z^p)_{p\ge 0}$ is a normal family. From every subsequence of $(\int g\, d\pi_z^p)_p$ one can thus extract a converging subsequence $(\int g\, d\pi_z^M)_M$ where $M = M(p)$ such that

$$\forall K \subset \mathbb{C}^+,\ K \text{ compact} \quad \sup_{z\in K} \left| \int g\, d\pi_z^M - \Gamma(g)(z) \right| \xrightarrow[M\to\infty]{} 0,$$

where $\Gamma(g)(z)$ is analytic over $\mathbb{C}^+$. If $z \in \mathbb{C}^\nabla$ and $\mathrm{Im}(z)$ is large enough, we know that $\int g\, d\pi_z^p \to \int g\, d\pi_z$. Therefore, $\Gamma(g)(z) = \int g\, d\pi_z$ for $z \in \mathbb{C}^\nabla$ and $\mathrm{Im}(z)$ large enough.

From this we can conclude that for all $z \in \mathbb{C}^+$, every subsequence has the same limit, say $\Gamma(g)(z)$ thus:

$$\forall z \in \mathbb{C}^+, \quad \int g \, d\pi_z^p \xrightarrow[p\to\infty]{} \Gamma(g)(z), \tag{3.4}$$

where $\Gamma(f)(z)$ is analytic. Moreover, it is straightforward to prove that

1. $\Gamma(a\, g + b\, h)(z) = a\Gamma(g) + b\Gamma(h)$,
2. $|\Gamma(g)(z)| \le \frac{\|g\|_\infty}{\mathrm{Im}(z)}$,
3. $\mathrm{Im}(\Gamma(g)(z)) \ge 0$ if $g \ge 0$ and $z \in \mathbb{C}^+$,
4. $\mathrm{Im}(z\Gamma(g)(z)) \ge 0$ if $g \ge 0$ and $z \in \mathbb{C}^+$.

As $\mathcal{H}$ is compact and since the application $g \mapsto \Gamma(g)(z)$ defined for $g \in C(\mathcal{H})$ is linear (property 1) and continuous (property 2), the Riesz representation theorem yields the existence of a measure $\pi_z(du, d\lambda) = \pi(z, du, d\lambda)$ such that

$$\Gamma(g)(z) = \int g(u, \lambda)\pi(z, du, d\lambda).$$

Similarly one can prove that

$$\int g \, d\tilde{\pi}_z^p \to \tilde{\Gamma}(g)(z) = \int g \, d\tilde{\pi}_z. \tag{3.5}$$

Let us now prove that π and $\tilde{\pi}$ satisfy (2.6) and (2.7). We first check that[1]

$$\forall z \in \mathbb{C}^+, \forall u \in [0, 1], \quad d(\pi_z)(u) \neq 0. \tag{3.6}$$

Indeed assume that for a given u, there exists $z_0 \in \mathbb{C}^+$ such that $d(\pi_{z_0})(u) = 0$ and consider the function $\Phi(z) = \int \sigma^2(t, cu)\, \pi(z, dt, d\zeta)$. As $d(\pi_z)(u) = 1 + \Phi(z)$, we have $\mathrm{Im}(\Phi(z_0)) = 0$. Since $\mathrm{Im}(\Phi(z))$ is harmonic and non-negative over $\mathbb{C}^+$, the mean value property implies that $\mathrm{Im}(\Phi(z)) = 0$ over $\mathbb{C}^+$. By the Cauchy-Riemann equations, $\mathrm{Re}(\Phi(z))$ is therefore constant. But since $|\Phi(z)| \le \frac{\sigma^2_{\max}}{\mathrm{Im}(z)}$, we have $\Phi(z) = 0$. This yields in particular $d(\pi_{z_0})(u) = 1$ which contradicts $d(\pi_{z_0})(u) = 0$.

Due to (3.4), (3.5) and (3.6), one has $\frac{1}{D(\tilde{\pi}_z^p, \pi_z^p)(u,\lambda)} \to_p \frac{1}{D(\tilde{\pi}_z, \pi_z)(u,\lambda)}$. Since $\frac{1}{D(\tilde{\pi}_z^p, \pi_z^p)(u,\lambda)} \le \frac{1}{\mathrm{Im}(z)}$, the dominated convergence theorem yields:

$$\int \frac{g(u, \lambda)}{D(\tilde{\pi}_z^p, \pi_z^p)(u, \lambda)} H(du, d\lambda) \xrightarrow[p\to\infty]{} \int \frac{g(u, \lambda)}{D(\tilde{\pi}_z, \pi_z)(u, \lambda)} H(du, d\lambda).$$

On the other hand $\int g \, d\pi_z^p \to_p \int g \, d\pi_z$ and (2.6) is established. One can establish Eq. (2.7) similarly.

It remains to prove that $f(z) = \int d\pi_z$ is the Stieltjes transform of a probability measure (one will prove similarly the corresponding result for $\tilde{f}(z) = \int d\tilde{\pi}_z$). Recall that

$$\mathrm{Im}(f(z)) = \mathrm{Im}\left(\int d\pi_z\right) \ge \int \frac{\mathrm{Im}(z)}{|D(\tilde{\pi}_z, \pi_z)(u, \lambda)|^2} H(du, d\lambda) > 0$$

by (2.6). Moreover, since $|f(z)| \le \frac{1}{\mathrm{Im}(z)}$, $f(z)$ is the Stieltjes transform of a subprobability measure. It remains to check that $\lim_{y\to+\infty} iyf(iy) = -1$. Since

$$\left| \int \sigma^2(u, t)\tilde{\pi}(iy, dt, d\zeta) \right| \le \frac{\sigma^2_{\max}}{y} \quad \text{and} \quad \left| \int \sigma^2(t, cu)\, \pi(iy, dt, d\zeta) \right| \le \frac{\sigma^2_{\max}}{y},$$

---

[1] Il existe peut-être un argument plus naturel basé sur $|d(\pi_z^p)| \ge \frac{\sqrt{2}}{2}$. A voir...

and

$$iyf(iy) = \int \frac{iy\, H(du, d\lambda)}{-iy(1 + \int \sigma^2(u,t)\tilde{\pi}(iy, dt, d\zeta)) + \frac{\lambda}{1+c\int \sigma^2(t,cu)\pi(iy,dt,d\zeta)}} \ ,$$

the Dominated convergence theorem yields the desired result. Theorem 2.3 is proved.

4. Proof of Theorem 2.4

We first give an outline of the proof. The proof is carried out following three steps:

(1) We first prove that for each subsequence $M(n)$ of n there exists a subsequence $M_{\rm sub} = M_{\rm sub}(n)$ such that for all $z \in \mathbb{C}^+$,

$$L_z^{M_{\rm sub}} \xrightarrow[n\to\infty]{w} \mu_z \quad \text{and} \quad \tilde{L}_z^{M_{\rm sub}} \xrightarrow[n\to\infty]{w} \tilde{\mu}_z, \tag{4.1}$$

where μ_z and $\tilde{\mu}_z$ are complex measures, a priori random, with support included in $\mathcal{H}$ (Section 4.1).

(2) We then prove that $z \mapsto \mu_z$ and $z \mapsto \tilde{\mu}_z$ are Stieltjes kernels (Section 4.2).

(3) We finally prove that for a countable collection of $z \in \mathbb{C}^+$ with a limit point, say

$$\mathcal{C} = \{z_p\}_{p\in\mathbb{N}} \cup \{z_\infty\} \quad \text{with} \quad z_p \to z_\infty,$$

the measures μ_z and $\tilde{\mu}_z$ (which are a priori random) satisfy equations (2.6) and (2.7) almost surely for all $z \in \mathcal{C}$. Since $\mathcal{C}$ has a limit point in $\mathbb{C}^+$, analyticity arguments will yield:

$$\text{a.s.} \quad \forall z \in \mathbb{C}^+, \quad \mu_z = \pi_z \quad \text{and} \quad \tilde{\mu}_z = \tilde{\pi}_z.$$

Otherwise stated,

$$\text{a.s.} \quad \forall z \in \mathbb{C}^+, \quad L_z^n \xrightarrow{w} \pi_z \quad \text{and} \quad \tilde{L}_z^n \xrightarrow{w} \tilde{\pi}_z$$

which yields the desired result (Section 4.3).

4.1. Step 1: convergence of subsequences $L_z^{M_{\rm sub}}$ and $\tilde{L}_z^{M_{\rm sub}}$.

Let $z_0 \in \mathbb{C}^+$ and let $B = \{z \in \mathbb{C}, |z - z_0| < \delta\} \subset \mathbb{C}^+$. Due to assumption (A-3) and to the fact that $|q_{ii}(z)| \le \mathrm{Im}^{-1}(z)$, Helly's theorem implies that for each subsequence of n there exists a subsequence $M = M(n)$ and a complex measure μ_{z_0} such that

$$L_{z_0}^M \xrightarrow[n\to\infty]{w} \mu_{z_0}.$$

Since L^n is random, μ_{z_0} is a priori random too but due to (A-3), its support is included in $\mathcal{H}$. Let $(z_k, k \ge 1)$ be a sequence of complex numbers dense in $\mathbb{C}^+$, then by Cantor diagonalization argument, one can extract a subsequence from M, say $M_{\rm sub}$, such that

$$\forall k \in \mathbb{N}, \quad L_{z_k}^{M_{\rm sub}} \xrightarrow[n\to\infty]{w} \mu_{z_k} \quad \text{and} \quad \tilde{L}_{z_k}^{M_{\rm sub}} \xrightarrow[n\to\infty]{w} \tilde{\mu}_{z_k},$$

where μ_{z_k} and $\tilde{\mu}_{z_k}$ are complex measures, a priori random. Let $g \in C(\mathcal{K})$ and let $z \in \mathbb{C}^+$. There exists z_k such that $|z - z_k| \le \epsilon$ and

$$\left| \int g \, dL_z^{M_{\mathrm{sub}}(n)} - \int g \, dL_z^{M_{\mathrm{sub}}(m)} \right| \le \overbrace{\left| \int g \, dL_z^{M_{\mathrm{sub}}(n)} - \int g \, dL_{z_k}^{M_{\mathrm{sub}}(n)} \right|}^{(a)} \\ + \underbrace{\left| \int g \, dL_{z_k}^{M_{\mathrm{sub}}(n)} - \int g \, dL_{z_k}^{M_{\mathrm{sub}}(m)} \right|}_{(b)} + \underbrace{\left| \int g \, dL_{z_k}^{M_{\mathrm{sub}}(m)} - \int g \, dL_z^{M_{\mathrm{sub}}(m)} \right|}_{(c)}$$

Let n and m be large enough. Since $L_{z_k}^{M_{\mathrm{sub}}}$ converges, (b) goes to zero. Since $q_{ii}(z)$ is analytic and since $|q_{ii}(z)| \le \mathrm{Im}^{-1}(z)$, there exists $K > 0$, such that

$$\forall i \ge 1, \ \forall z, z' \text{ close enough}, \quad |q_{ii}(z) - q_{ii}(z')| \le K|z - z'|.$$

Thus $\max\{(a), (c)\} \le K\|g\|_\infty |z - z_k|$. Therefore, $\left(\int g \, dL_z^{M_{\mathrm{sub}}}\right)$ is a Cauchy sequence and converges to $\Theta(g)(z)$. Since $g \mapsto \Theta(g)(z)$ is linear and since $|\Theta(g)(z)| \le \mathrm{Im}^{-1}(z)\|g\|_\infty$, Riesz representation's theorem yields the existence of μ_z such that $\Theta(g)(z) = \int g \, d\mu_z$ (recall that the support of μ_z is included in $\mathcal{H}$ which is compact). The convergence of $\tilde{L}_z^{M_{\mathrm{sub}}}$ can be proved similarly and (4.1) is satisfied. The first step is proved.

4.2. Step 2: the kernels μ_z and $\tilde{\mu}_z$ are Stieltjes kernels.

Let us now prove that $z \mapsto \int g \, d\mu_z$ is analytic over $\mathbb{C}^+$. Since $\left|\int g \, dL_z^{M_{\mathrm{sub}}}\right| \le \mathrm{Im}^{-1}(z)\|g\|_\infty$, from each subsequence of $\left(\int g \, dL_z^{M_{\mathrm{sub}}}\right)$, one can extract a subsequence that converges to an analytic function. Since this limit is equal to $\int g \, d\mu_z$, the analyticity of $z \mapsto \int g \, d\mu_z$ over $\mathbb{C}^+$ is proved. Since properties (3) and (4) defining the Stieltjes kernels are satisfied by L_z^n, the kernel μ_z inherits them. Therefore, μ_z is a Stieltjes kernel. Similarly, one can prove that $\tilde{\mu}_z$ is a Stieltjes kernel. The second step is proved.

4.3. Step 3: the kernels μ_z and $\tilde{\mu}_z$ are almost surely equal to π_z and $\tilde{\pi}_z$.

We will now prove that almost surely μ_z and $\tilde{\mu}_z$ satisfy equations (2.6) and (2.7).

In the sequel we will drop the subscript n from the notations relative to matrices, and the superscript n from Λ_{ii}^n. Let $\vec{e}_i = (\delta_i(k))_{1 \le k \le n}$ and $\vec{f}_i = (\delta_i(k))_{1 \le k \le N}$. For the sake of simplicity, Σ^T will be denoted Ξ. Consider the following notations:

Matrix	Y	Λ	Σ	$\Sigma^T_{(i)}$	Y^T	Λ^T	Ξ	$\Xi^T_{(i)}$
ith row	$\vec{y}_{i\cdot}$	$\Lambda_{ii}\vec{e}_i$	$\vec{\xi}_{i\cdot}$	$\vec{\eta}_{i\cdot}$	$\vec{y}_{\cdot i}$	$\Lambda_{ii}\vec{f}_i$	$\vec{\xi}_{\cdot i}$	$\vec{\eta}_{\cdot i}$
Matrix when ith row is deleted	-	-	$\Sigma_{(i)}$	$\Sigma^T_{(i,i)}$	-	-	$\Xi_{(i)}$	$\Xi^T_{(i,i)}$

In particular, $\vec{\xi}_{i\cdot} = \vec{y}_{i\cdot} + \Lambda_{ii}\vec{e}_i$ and $\vec{\xi}_{\cdot i} = \vec{y}_{\cdot i} + \Lambda_{ii}\vec{f}_i$ for $1 \le i \le N$. We will denote by D_i and Δ_j the respectively $n \times n$ and $N \times N$ diagonal matrices defined by

$$D_i = \mathrm{diag}\left(\frac{\sigma\left(\frac{i}{N}, \frac{1}{n}\right)}{\sqrt{n}}, \cdots, \frac{\sigma\left(\frac{i}{N}, 1\right)}{\sqrt{n}}\right), \quad \Delta_j = \mathrm{diag}\left(\frac{\sigma\left(\frac{1}{N}, \frac{j}{n}\right)}{\sqrt{n}}, \cdots, \frac{\sigma\left(1, \frac{j}{n}\right)}{\sqrt{n}}\right).$$

Finally, for $1 \le i \le N$, we denote by $D_{(i,i)}$ and $\Delta_{(i,i)}$ the matrices that remain after deleting row i and column i from D_i and Δ_i respectively.

We can state our first lemma:

Lemma 4.1. *Let $z \in \mathbb{C}^+$ be fixed.*

(1) *The ith diagonal element $q_{ii}(z)$ of the matrix $(\Sigma\Sigma^T - zI_N)^{-1}$ can be written :*

$$q_{ii}(z) = \frac{1}{-z - \frac{z}{n}\sum_{k=1}^{n} \sigma^2\left(\frac{i}{N}, \frac{k}{n}\right)\tilde{q}_{kk}(z) + \frac{\Lambda_{ii}^2}{1+\frac{1}{n}\sum_{k=1}^{N}\sigma^2\left(\frac{k}{N},\frac{i}{n}\right)q_{ii}(z)+\varepsilon_{i,n}^{(4)}+\varepsilon_{i,n}^{(5)}} + \Lambda_{ii}\,\varepsilon_{i,n}^{(1)} + \varepsilon_{i,n}^{(2)} + \varepsilon_{i,n}^{(3)}} \tag{4.2}$$

where $1 \le i \le N$ and

$$\begin{aligned}
\varepsilon_{i,n}^{(1)} &= -z\,\vec{y}_{i\cdot}\left(\Sigma_{(i)}^T\Sigma_{(i)} - zI\right)^{-1}\vec{e}_i^{\,T} - z\,\vec{e}_i\left(\Sigma_{(i)}^T\Sigma_{(i)} - zI\right)^{-1}\vec{y}_{i\cdot}^{\,T}\\
\varepsilon_{i,n}^{(2)} &= -z\,\vec{y}_{i\cdot}\left(\Sigma_{(i)}^T\Sigma_{(i)} - zI\right)^{-1}\vec{y}_{i\cdot}^{\,T} + z\,\mathrm{Tr}\left(D_i^2\left(\Sigma_{(i)}^T\Sigma_{(i)} - zI\right)^{-1}\right)\\
\varepsilon_{i,n}^{(3)} &= z\,\mathrm{Tr}\left(D_i^2\left(\Sigma^T\Sigma - zI\right)^{-1}\right) - z\,\mathrm{Tr}\left(D_i^2\left(\Sigma_{(i)}^T\Sigma_{(i)} - zI\right)^{-1}\right)\\
\varepsilon_{i,n}^{(4)} &= \vec{\eta}_{i\cdot}\left(\Sigma_{(i,i)}\Sigma_{(i,i)}^T - zI\right)^{-1}\vec{\eta}_{i\cdot}^{\,T} - \mathrm{Tr}\left(\Delta_{(i,i)}^2\left(\Sigma_{(i,i)}\Sigma_{(i,i)}^T - zI\right)^{-1}\right)\\
\varepsilon_{i,n}^{(5)} &= \mathrm{Tr}\left(\Delta_{(i,i)}^2\left(\Sigma_{(i,i)}\Sigma_{(i,i)}^T - zI\right)^{-1}\right) - \mathrm{Tr}\left(\Delta_i^2\left(\Sigma\Sigma^T - zI\right)^{-1}\right)
\end{aligned}$$

Moreover almost surely,

$$\forall k, 1 \le k \le 5, \quad \lim_{n\to\infty}\frac{1}{N}\sum_{i=1}^{N}\left|\varepsilon_{i,n}^{(k)}\right| = 0. \tag{4.3}$$

(2) *If $1 \le i \le N$ then the ith diagonal element $\tilde{q}_{ii}(z)$ of the matrix $(\Sigma^T\Sigma - zI_n)^{-1} = (\Xi\Xi^T - zI_n)^{-1}$ can be written:*

$$\tilde{q}_{ii}(z) = \frac{1}{-z - \frac{z}{n}\sum_{k=1}^{N} \sigma^2\left(\frac{k}{N}, \frac{i}{n}\right)q_{kk}(z) + \frac{\Lambda_{ii}^2}{1+\frac{1}{n}\sum_{k=1}^{n}\sigma^2\left(\frac{i}{N},\frac{k}{n}\right)\tilde{q}_{ii}(z)+\tilde{\varepsilon}_{i,n}^{(4)}+\tilde{\varepsilon}_{i,n}^{(5)}} + \Lambda_{ii}\,\tilde{\varepsilon}_{i,n}^{(1)} + \tilde{\varepsilon}_{i,n}^{(2)} + \tilde{\varepsilon}_{i,n}^{(3)}}. \tag{4.4}$$

If $N+1 \le i \le n$, then $\tilde{q}_{ii}$ can be written:

$$\tilde{q}_{ii}(z) = \frac{1}{-z - \frac{z}{n}\sum_{k=1}^{N} \sigma^2\left(\frac{k}{N}, \frac{i}{n}\right)q_{kk}(z) + \tilde{\varepsilon}_{i,n}^{(2)} + \tilde{\varepsilon}_{i,n}^{(3)}}. \tag{4.5}$$

where

$$\begin{aligned}
\tilde{\varepsilon}_{i,n}^{(1)} &= -z\,\vec{y}_{\cdot i}\left(\Xi_{(i)}^T\Xi_{(i)} - zI\right)^{-1}\vec{f}_i^{\,T} - z\,\vec{f}_i\left(\Xi_{(i)}^T\Xi_{(i)} - zI\right)^{-1}\vec{y}_{\cdot i}^{\,T}\\
\tilde{\varepsilon}_{i,n}^{(2)} &= -z\,\vec{y}_{\cdot i}\left(\Xi_{(i)}^T\Xi_{(i)} - zI\right)^{-1}\vec{y}_{\cdot i}^{\,T} + z\,\mathrm{Tr}\left(\Delta_i^2\left(\Xi_{(i)}^T\Xi_{(i)} - zI\right)^{-1}\right)\\
\tilde{\varepsilon}_{i,n}^{(3)} &= z\,\mathrm{Tr}\left(\Delta_i^2\left(\Xi^T\Xi - zI\right)^{-1}\right) - z\,\mathrm{Tr}\left(\Delta_i^2\left(\Xi_{(i)}^T\Xi_{(i)} - zI\right)^{-1}\right)\\
\tilde{\varepsilon}_{i,n}^{(4)} &= \vec{\eta}_{\cdot i}\left(\Xi_{(i,i)}\Xi_{(i,i)}^T - zI\right)^{-1}\vec{\eta}_{\cdot i}^{\,T} - \mathrm{Tr}\left(D_{(i,i)}^2\left(\Xi_{(i,i)}\Xi_{(i,i)}^T - zI\right)^{-1}\right)\\
\tilde{\varepsilon}_{i,n}^{(5)} &= \mathrm{Tr}\left(D_{(i,i)}^2\left(\Xi_{(i,i)}\Xi_{(i,i)}^T - zI\right)^{-1}\right) - \mathrm{Tr}\left(D_i^2\left(\Xi\Xi^T - zI\right)^{-1}\right)
\end{aligned}$$

Moreover, almost surely

$$\begin{aligned} \textit{for } k &= 1,4,5 \quad \lim_{n\to\infty} \frac{1}{N}\sum_{i=1}^{N} \left|\tilde{\varepsilon}_{i,n}^{(k)}\right| = 0, \\ \textit{for } k &= 2,3 \quad \lim_{n\to\infty} \frac{1}{n}\sum_{i=1}^{n} \left|\tilde{\varepsilon}_{i,n}^{(k)}\right| = 0. \end{aligned} \tag{4.6}$$

Proof of Lemma 4.1. Since $q_{ii}(z) = (\Sigma\Sigma^T - zI)^{-1}_{ii}$, this element is the inverse of the Schur complement of $\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)$ in $\left(\Sigma\Sigma^T - zI\right)$ (see for instance [11], Appendix A). In other words

$$q_{ii}(z) = \left(\|\vec{\xi}_{i\cdot}\|^2 - z - \vec{\xi}_{i\cdot}\Sigma_{(i)}^T(\Sigma_{(i)}\Sigma_{(i)}^T - zI)^{-1}\Sigma_{(i)}\vec{\xi}_{i\cdot}^T\right)^{-1} .$$

Using the identity

$$I - \Sigma_{(i)}^T(\Sigma_{(i)}\Sigma_{(i)}^T - zI)^{-1}\Sigma_{(i)} = -z(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1} ,$$

we have

$$\begin{aligned} q_{ii}(z) &= \frac{1}{-z - z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T} \\ &= \frac{1}{-z - z\vec{y}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{y}_{i\cdot}^T - z\Lambda_{ii}^2\vec{e}_i(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{e}_i^T + \Lambda_{ii}\varepsilon_{i,n}^{(1)}} \\ &= \frac{1}{-z - \frac{z}{n}\sum_{k=1}^n \sigma^2\left(\frac{i}{N},\frac{k}{n}\right)\tilde{q}_{kk}(z) - z\Lambda_{ii}^2\vec{e}_i(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{e}_i^T + \Lambda_{ii}\varepsilon_{i,n}^{(1)} + \varepsilon_{i,n}^{(2)} + \varepsilon_{i,n}^{(3)}}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \vec{e}_i(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{e}_i^T &= (\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}_{ii} \\ &= \frac{1}{-z - z\vec{\eta}_{i\cdot}(\Sigma_{(ii)}\Sigma_{(ii)}^T - zI)^{-1}\vec{\eta}_{i\cdot}^T} \\ &= \frac{1}{-z\left(1 + \frac{1}{n}\sum_{k=1}^N \sigma^2\left(\frac{k}{N},\frac{i}{n}\right)q_{kk}(z) + \varepsilon_{i,n}^{(4)} + \varepsilon_{i,n}^{(5)}\right)} \end{aligned} \tag{4.7}$$

And (4.2) is established. It is important to already note that since $\vec{\eta}_{i\cdot}$ is the ith row of $\Sigma_{(i)}^T$, $\mathbb{E}\vec{\eta}_{i\cdot} = 0$ (while $\mathbb{E}\vec{y}_{i\cdot} = (0,\cdots,\Lambda_{ii},\cdots,0)$). If $i \le N$, (4.4) can be established in the same way. If $i \ge N+1$, then $\vec{\xi}_{\cdot i}$ is centered: There are no more Λ_{ii} and all the terms involving Λ_{ii} disappear in (4.4), which yields (4.5).

We now prove that

$$\frac{1}{N}\sum_{i=1}^{N}\left|\varepsilon_{i,n}^{(1)}\right| \xrightarrow[n\to\infty]{\text{a.s.}} 0. \tag{4.8}$$

One will prove similarly that $\frac{1}{N}\sum_{i=1}^N |\tilde{\varepsilon}_{i,n}^{(1)}| \to 0$ a.s. Denote by $R_n = (\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1} = (\rho_{ij})$. Since R_n is symetric, $\varepsilon_{i,n}^{(1)} = -z2\vec{y}_{i\cdot}R_n\vec{e}_i^T$ and

$$|\vec{y}_{i\cdot}R_n\vec{e}_i^T|^4 = |\sum_{k=1}^n Y_{ik}\rho_{ki}|^4 = \sum_{k_1,k_2,l_1,l_2} Y_{ik_1}Y_{ik_2}Y_{il_1}Y_{il_2}\rho_{k_1 i}\,\bar{\rho}_{k_2 i}\,\rho_{l_1 i}\,\bar{\rho}_{l_2 i}.$$

Denote by $\mathbb{E}_{R_n}$ the expectation conditionnally to the σ-algebra generated by R_n. Since $\vec{y}_{i\cdot}$ and R_n are independent and since $\mathbb{E}Y_{ik} = 0$, we get:

$$\begin{aligned}
\mathbb{E}_{R_n}|\vec{y}_{i\cdot}R_n\vec{e}_i^T|^4 &= 2\mathbb{E}_{R_n}\sum_{k,l;k\neq l} Y_{ik}^2|\rho_{ki}|^2 Y_{il}^2|\rho_{li}|^2 \\
&\quad +\mathbb{E}_{R_n}\sum_{k,l;k\neq l} Y_{ik}^2\rho_{ki}^2 Y_{il}^2\bar{\rho}_{li}^2 + \mathbb{E}_{R_n}\sum_k Y_{ik}^4|\rho_{ki}|^4 \\
&\leq 4\mathbb{E}(X_{ij}^n)^4\frac{\sigma_{\max}^4}{n^2}\sum_{k,l}|\rho_{ki}|^2|\rho_{li}|^2 = 4\mathbb{E}(X_{ij}^n)^4\frac{\sigma_{\max}^4}{n^2}\left(\sum_k|\rho_{ki}|^2\right)^2
\end{aligned}$$

but $\sum_k|\rho_{ki}|^2 = \|R_n e_i\|^2 \leq \|R_n\|^2 \leq \frac{1}{\mathrm{Im}^2(z)}$. Therefore,

$$\mathbb{E}\left|\varepsilon_{i,n}^{(1)}\right|^4 \leq \frac{|2z|^4 4\mathbb{E}(X_{ij}^n)^4\sigma_{\max}^4}{n^2\mathrm{Im}^4(z)} \propto \frac{1}{n^2}. \tag{4.9}$$

Finally,

$$\begin{aligned}
\mathbb{P}\left\{\frac{1}{N}\sum_{i=1}^N\left|\varepsilon_{i,n}^{(1)}\right| > \delta\right\} &\leq \frac{1}{\delta^4N^4}\mathbb{E}\left(\sum_{i=1}^N|\varepsilon_{i,n}^{(1)}|\right)^4 \\
&\overset{(a)}{\leq} \frac{1}{\delta^4N^4}\left(\sum_{i=1}^N\left(\mathbb{E}|\varepsilon_{i,n}^{(1)}|^4\right)^{\frac{1}{4}}\right)^4 \leq \frac{1}{\delta^4}\sup_{1\leq i\leq N}\mathbb{E}|\varepsilon_{i,n}^{(1)}|^4 \overset{(b)}{\propto} \frac{1}{n^2}
\end{aligned}$$

where (a) follows from Minkowski's inequality and (b) from (4.9) and Borel-Cantelli's lemma yields Eq. (4.8).

Let us now prove that

$$\frac{1}{N}\sum_{i=1}^N\left|\varepsilon_{i,n}^{(2)}\right| \xrightarrow[n\to\infty]{\text{a.s.}} 0. \tag{4.10}$$

One will prove similarly that $\frac{1}{n}\sum|\tilde{\varepsilon}_{i,n}^{(2)}|$, $\frac{1}{N}\sum|\varepsilon_{i,n}^{(4)}|$ and $\frac{1}{N}\sum|\tilde{\varepsilon}_{i,n}^{(4)}|$ go to zero a.s. Denote by $\vec{x}_{i\cdot} = (X_{i1},\cdots,X_{in})$ and write $\vec{y}_{i\cdot} = \vec{x}_{i\cdot}D_i$. In particular,

$$\vec{y}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{y}_{i\cdot}^T = \vec{x}_{i\cdot}D_i(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}D_i^T\vec{x}_{i\cdot}^T$$

where $\vec{x}_{i\cdot}$ and $D_i(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}D_i^T$ are independent. Lemma 2.7 in [3] states that

$$\begin{aligned}
&\mathbb{E}|\vec{x}_{i\cdot}C\vec{x}_{i\cdot}^T - \mathrm{Tr}\,C|^p \\
&\qquad\leq K_p\left(\left(\mathbb{E}(X_{i1})^4\mathrm{Tr}\,CC^T\right)^{p/2} + \mathbb{E}(X_{i1})^{2p}\mathrm{Tr}(CC^T)^{p/2}\right)
\end{aligned} \tag{4.11}$$

for all $p \geq 2$. Take $p = 2 + \epsilon/2$ where ϵ is given by (A-1) and let $C = D_i(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}D_i^T$. Then

$$\forall q \geq 1, \quad \mathrm{Tr}(CC^T)^q \leq \frac{\sigma_{\max}^{4q}}{n^{2q-1}} \times \frac{1}{\mathrm{Im}^{2q}(z)}. \tag{4.12}$$

Therefore, (4.11) and (4.12) yield

$$\mathbb{E}|\vec{x}_{i\cdot}C\vec{x}_{i\cdot}^T - \mathrm{Tr}\,C|^{2+\epsilon/2} \leq \frac{K_1}{n^{1+\epsilon/4}} + \frac{K_2}{n^{1+\epsilon}} \leq \frac{K}{n^{1+\epsilon/4}}$$

where the constants K, K_1 and K_2 depend on the moments of X_{i1}, on $\sigma_{\max}$ and on $\mathrm{Im}(z)$. Thus

$$\mathbb{E}\left|\varepsilon_{i,n}^{(2)}\right|^p \leq \frac{K|z|^p}{n^{1+\epsilon/4}}. \tag{4.13}$$

Finally,

$$\begin{aligned}\mathbb{P}\left\{\frac{1}{N}\sum_{i=1}^{N}\left|\varepsilon_{i,n}^{(2)}\right|>\delta\right\} &\leq \frac{1}{\delta^p N^p}\mathbb{E}\left(\sum_{i=1}^{N}|\varepsilon_{i,n}^{(2)}|\right)^p\\ &\overset{(a)}{\leq} \frac{1}{\delta^p N^p}\left(\sum_{i=1}^{N}\left(\mathbb{E}|\varepsilon_{i,n}^{(2)}|^p\right)^{\frac{1}{p}}\right)^p\\ &\leq \frac{1}{\delta^p}\sup_{1\leq i\leq N}\mathbb{E}|\varepsilon_{i,n}^{(2)}|^p \overset{(b)}{\propto} \frac{1}{n^{1+\epsilon/4}}\end{aligned}$$

where (a) follows from Minkowski's inequality and (b) from (4.13), and Borel-Cantelli's lemma yields (4.10).
We now prove that

$$\frac{1}{N}\sum_{i=1}^{N}\left|\varepsilon_{i,n}^{(3)}\right| \xrightarrow[n\to\infty]{\text{a.s.}} 0. \tag{4.14}$$

One will prove similarly that $\frac{1}{n}\sum|\tilde{\varepsilon}_{i,n}^{(3)}|$ goes to zero. Since $\Sigma^T\Sigma=\Sigma_{(i)}^T\Sigma_{(i)}+\vec{\xi}_{i\cdot}^T\vec{\xi}_{i\cdot}$, Lemma 2.6 in [16] yields:

$$\left|\mathrm{Tr}\left((\Sigma^T\Sigma-zI)^{-1}-(\Sigma_{(i)}^T\Sigma_{(i)}-zI)^{-1}\right)D_i^2\right|\leq\frac{\sigma_{\max}^2}{n\mathrm{Im}(z)},$$

In particular,

$$\left|\varepsilon_{i,n}^{(3)}\right|\leq\frac{|z|\sigma_{\max}^2}{n\mathrm{Im}(z)} \tag{4.15}$$

which immediatly yields (4.14).
We finally prove that

$$\frac{1}{N}\sum_{i=1}^{N}\left|\varepsilon_{i,n}^{(5)}\right| \xrightarrow[n\to\infty]{\text{a.s.}} 0. \tag{4.16}$$

One will prove similarly that $\frac{1}{N}\sum|\tilde{\varepsilon}_{i,n}^{(5)}|$ goes to zero. Write

$$\begin{aligned}\varepsilon_{i,n}^{(5)}=\mathrm{Tr}\,\Delta_{(i,i)}^2\left(\Sigma_{(i,i)}\Sigma_{(i,i)}^T-zI\right)^{-1}-\mathrm{Tr}\,\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T-zI\right)^{-1}\\ +\mathrm{Tr}\,\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T-zI\right)^{-1}-\mathrm{Tr}\,\Delta_i^2\left(\Sigma\Sigma^T-zI\right)^{-1}\end{aligned}$$

As for $\varepsilon_{i,n}^{(3)}$, one can prove that

$$\left|\mathrm{Tr}\,\Delta_{(i,i)}^2\left(\Sigma_{(i,i)}\Sigma_{(i,i)}^T-zI\right)^{-1}-\mathrm{Tr}\,\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T-zI\right)^{-1}\right|\leq\frac{\sigma_{\max}^2}{n\mathrm{Im}(z)}$$

by applying Lemma 2.6 in [16]. Let

$$\kappa_{i,n}=\mathrm{Tr}\,\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T-zI\right)^{-1}-\mathrm{Tr}\,\Delta_i^2\left(\Sigma\Sigma^T-zI\right)^{-1}.$$

By applying to $\Sigma\Sigma^T - zI$ the identities relative to the inverse of a partitioned matrix (see [11], Appendix A), we obtain: $\operatorname{Tr}\Delta_i^2\left(\Sigma\Sigma^T - zI\right)^{-1} = \Psi_1 + \Psi_2 + \Psi_3$ where

$$\begin{aligned}
\Psi_1 &= \operatorname{Tr}\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}\\
\Psi_2 &= \frac{\operatorname{Tr}\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}\Sigma_{(i)}\vec{\xi}_{i\cdot}^T\vec{\xi}_{i\cdot}\Sigma_{(i)}^T\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}}{-z - z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T}\\
\Psi_3 &= \frac{1}{n}\frac{\sigma^2\left(\frac{i}{N},\frac{i}{n}\right)}{-z - z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T}\ .
\end{aligned}$$

In particular, $\kappa_{i,n} = -\Psi_2 - \Psi_3$. We have

$$\begin{aligned}
|\Psi_2| &= \left|\frac{\vec{\xi}_{i\cdot}\Sigma_{(i)}^T\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}\Delta_{(i,i)}^2\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}\Sigma_{(i)}\vec{\xi}_{i\cdot}^T}{-z - z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T}\right|\\
&\leq \left\|\Delta_{(i,i)}\right\|^2\frac{\left\|\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}\Sigma_{(i)}\vec{\xi}_{i\cdot}^T\right\|^2}{\left|z + z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T\right|}
\end{aligned}$$

Let $\Sigma_{(i)} = \sum_{l=1}^{N-1}\nu_l u_l v_l^T$ be a singular value decomposition of $\Sigma_{(i)}$ where ν_l, u_l, and v_l are respectively the singular values, left singular vectors, and right singular vectors of $\Sigma_{(i)}$. Then

$$\left\|\left(\Sigma_{(i)}\Sigma_{(i)}^T - zI\right)^{-1}\Sigma_{(i)}\vec{\xi}_{i\cdot}^T\right\|^2 = \sum_{l=1}^{N-1}\frac{\nu_l^2\left|v_l^T\vec{\xi}_{i\cdot}^T\right|^2}{\left|\nu_l^2 - z\right|^2}$$

and

$$\operatorname{Im}\left(z + z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T\right) = \operatorname{Im}(z)\left(1 + \sum_{l=1}^{N-1}\frac{\nu_l^2\left|v_l^T\vec{\xi}_{i\cdot}^T\right|^2}{\left|\nu_l^2 - z\right|^2}\right)\ .$$

As a consequence, $|\Psi_2| \leq \frac{\sigma_{\max}^2}{n}\frac{1}{\operatorname{Im}(z)}$. Furthermore, since $\operatorname{Im}\left(z\vec{\xi}_{i\cdot}(\Sigma_{(i)}^T\Sigma_{(i)} - zI)^{-1}\vec{\xi}_{i\cdot}^T\right) \geq 0$ by Proposition 2.2-(3), we have $|\Psi_3| \leq \frac{\sigma_{\max}^2}{n}\frac{1}{\operatorname{Im}(z)}$. Thus, $\left|\varepsilon_{i,n}^{(5)}\right| \leq \frac{\sigma_{\max}^2}{n}\frac{3}{\operatorname{Im}(z)}$, which immediatly yields (4.16). Lemma 4.1 is proved. □

Recall notation D introduced at the beginning of Section 3:

$$D(\tilde{\pi}_z, \pi_z)(u,\lambda) = -z\left(1 + \int\sigma^2(u,t)\tilde{\pi}(z,dt,d\zeta)\right) + \frac{\lambda}{1 + c\int\sigma^2(t,cu)\pi(z,dt,d\zeta)}.$$

Corollary 4.2. *Let $z \in \mathbb{C}^+$ be fixed. Then almost surely*

$$\forall g \in C(\mathcal{K}),\qquad \lim_{n\to\infty}\left|\frac{1}{N}\sum_{i=1}^N q_{ii}^n g(\Lambda_{ii}^2, i/N) - \frac{1}{N}\sum_{i=1}^N\frac{g(\Lambda_{ii}^2, i/N)}{D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N}\right)}\right| = 0. \tag{4.17}$$

Proof of Corollary 4.2. Following the notations D and d introduced at the beginning of Section 3, we introduce their empirical counterparts:

$$\begin{aligned} d^n(u) &= 1+\frac{1}{n}\sum_{k=1}^{N}\sigma^2\left(\frac{k}{N},\frac{N}{n}u\right)q_{ii}(z)+\varepsilon_{i,n}^{(4)}+\varepsilon_{i,n}^{(5)} \\ D^n(u) &= -z-\frac{z}{n}\sum_{k=1}^{n}\sigma^2\left(u,\frac{k}{n}\right)\tilde{q}_{kk}(z)+\frac{\Lambda_{ii}^2}{d^n(u)} \\ &\quad +\Lambda_{ii}\,\varepsilon_{i,n}^{(1)}+\varepsilon_{i,n}^{(2)}+\varepsilon_{i,n}^{(3)} \end{aligned}$$

Since $q_{ii}=(D^n(i/N))^{-1}$ by (4.2) and $(\Sigma_{(i)}^T\Sigma_{(i)}-zI)_{ii}^{-1}=(-zd^n(i/N))^{-1}$ by (4.7), Proposition 2.2-(1) yields:

$$\frac{1}{|D^n(i/N)|}\le\frac{1}{\mathrm{Im}(z)}\quad\text{and}\quad\frac{1}{|d^n(i/N)|}\le\frac{|z|}{\mathrm{Im}(z)}. \tag{4.18}$$

On the other hand, since μ_z and $\tilde{\mu}_z$ are Stieltjes kernels, we have:

$$\frac{1}{|D(\tilde{\mu}_z,\mu_z)(i/N,\Lambda_{ii}^2)|}\le\frac{1}{\mathrm{Im}(z)}\quad\text{and}\quad\frac{1}{|d(\mu_z)(i/N)|}\le\frac{|z|}{\mathrm{Im}(z)}. \tag{4.19}$$

Therefore,

$$\begin{aligned} & q_{ii}^n-\frac{1}{D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N},\Lambda_{ii}^2\right)} = \frac{-z\left(\int\sigma^2(i/N,\cdot)\,d\tilde{L}_z^n-\int\sigma^2(i/N,\cdot)\,d\tilde{\mu}_z\right)}{D^n\left(\frac{i}{N}\right)\times D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N},\Lambda_{ii}^2\right)} \\ + \quad & \frac{\Lambda_{ii}\,\varepsilon_{i,n}^{(1)}+\varepsilon_{i,n}^{(2)}+\varepsilon_{i,n}^{(3)}}{D^n\left(\frac{i}{N}\right)\times D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N},\Lambda_{ii}^2\right)} \\ + \quad & \frac{\Lambda_{ii}^2\left(\frac{N}{n}\int\sigma^2(\cdot,i/n)\,dL_z^n-c\int\sigma^2(\cdot,i/n)\,d\mu_z\right)}{d(\mu_z)\left(\frac{i}{N}\right)\times d^n\left(\frac{i}{N}\right)\times D^n\left(\frac{i}{N}\right)\times D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N},\Lambda_{ii}^2\right)} \\ + \quad & \frac{\Lambda_{ii}^2\left(\varepsilon_{i,n}^{(4)}+\varepsilon_{i,n}^{(5)}\right)}{d(\mu_z)\left(\frac{i}{N}\right)\times d^n\left(\frac{i}{N}\right)\times D^n\left(\frac{i}{N}\right)\times D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N},\Lambda_{ii}^2\right)} \end{aligned}$$

Recall that the Λ_{ii}'s are assumed to be bounded (say $|\Lambda_{ii}|\le K$). Due to (4.18) and (4.19), we get:

$$\begin{aligned} & \left|q_{ii}^n-\frac{1}{D(\tilde{\mu}_z,\mu_z)\left(\frac{i}{N},\Lambda_{ii}^2\right)}\right| \le \frac{|z|}{\mathrm{Im}^2(z)}\underbrace{\left|\int\sigma^2(i/N,\cdot)\,d\tilde{L}_z^n-\int\sigma^2(i/N,\cdot)\,d\tilde{\mu}_z\right|}_{I(i,n)} \\ + \quad & \frac{K\left|\varepsilon_{i,n}^{(1)}\right|+\left|\varepsilon_{i,n}^{(2)}\right|+\left|\varepsilon_{i,n}^{(3)}\right|}{\mathrm{Im}^2(z)} \\ + \quad & \frac{|z|^2K^2}{\mathrm{Im}^4(z)}\left(\underbrace{\left|\frac{N}{n}\int\sigma^2(\cdot,i/n)\,dL_z^n-c\int\sigma^2(\cdot,i/n)\,d\mu_z\right|}_{J(i,n)}+\left|\varepsilon_{i,n}^{(4)}\right|+\left|\varepsilon_{i,n}^{(5)}\right|\right) \end{aligned}$$

In order to prove $\sup_{i\leq N} I(i,n) \to 0$, recall that $C([0,1]^2) = C([0,1]) \otimes C([0,1])$. In particular, $\forall \epsilon > 0$, there exists $k \in \mathbb{N}$, $g_l \in C([0,1])$ and $h_l \in C([0,1])$ for $l \leq k$ such that $\sup_{x,t} \left| \sigma^2(x,t) - \sum_{l=1}^k g_l(x) h_l(t) \right| \leq \epsilon$. Therefore,

$$\left| \int \sigma^2(i/N,\cdot)\, d\tilde{L}_z^n - \int \sigma^2(i/N,\cdot)\, d\tilde{\mu}_z \right| \leq \sup_x \left| \int \sigma^2(x,\cdot)\, d\tilde{L}_z^n - \int \sigma^2(x,\cdot)\, d\tilde{\mu}_z \right| \xrightarrow[n\to\infty]{} 0$$

which implies that $\sup_{i\leq N} |I(i,n)|$ goes to zero. One can prove similarly that $\sup_{i\leq N} J(i,n)$ goes to zero. Therefore,

$$\begin{aligned} &\left| \frac{1}{N} \sum_{i=1}^N q_{ii}^n g(\Lambda_{ii}^2, i/N) - \frac{1}{N} \sum_{i=1}^N \frac{g(\Lambda_{ii}^2, i/N)}{D(\tilde{\mu}_z, \mu_z)\left(\frac{i}{N}, \Lambda_{ii}\right)} \right| \\ &\quad \leq \frac{|z|\|g\|_\infty}{\mathrm{Im}^2(z)} \sup_{i\leq N} I(i,n) + \frac{\|g\|_\infty}{\mathrm{Im}^2(z)} \left(K \frac{1}{N} \sum_{i=1}^N |\varepsilon_{i,n}^{(1)}| + \frac{1}{N} \sum_{i=1}^N |\varepsilon_{i,n}^{(2)}| + \frac{1}{N} \sum_{i=1}^N |\varepsilon_{i,n}^{(3)}| \right) \\ &\qquad + \frac{|z|^2 K^2 \|g\|_\infty}{\mathrm{Im}^4(z)} \left(\sup_{i\leq N} J(i,n) + \frac{1}{N} \sum_{i=1}^N |\varepsilon_{i,n}^{(4)}| + \frac{1}{N} \sum_{i=1}^N |\varepsilon_{i,n}^{(5)}| \right). \end{aligned}$$

and (4.17) is proved with the help of Lemma 4.1. □

We now come back to the proof of the third step of Theorem 2.4. For simplicity, we will denote by $n^* = M_{\mathrm{sub}}(n)$ where M_{sub} is defined previously, by $N^* = N(n^*)$.

A direct application of the Dominated convergence theorem yields that $(\lambda, u) \mapsto \frac{g(\lambda,u)}{D(\tilde{\mu}_z,\mu_z)(u,\lambda)}$ is bounded and continuous therefore (A-3) yields

$$\frac{1}{N^*} \sum_{i=1}^{N^*} \frac{g(\Lambda_{ii}^2, i/N^*)}{D(\tilde{\mu}_z, \mu_z)(i/N^*, \Lambda_{ii}^2)} \xrightarrow[n\to\infty]{} \int \frac{g(\lambda,u)}{D(\tilde{\mu}_z,\mu_z)(u,\lambda)} H(d\lambda, du). \tag{4.20}$$

Moreover,

$$\frac{1}{N^*} \sum_{i=1}^{N^*} g(\Lambda_{ii}^2, i/N^*) q_{ii} \xrightarrow[n\to\infty]{} \int g\, d\mu_z. \tag{4.21}$$

Consider now a countable set $\mathcal{C}$ with a limit point. Since $\mathcal{C}$ is countable, (4.17) holds almost surely for every $z \in \mathcal{C}$ and for every $g \in C(\mathcal{K})$. Thus (4.20) and (4.21) yield that μ_z and $\tilde{\mu}_z$ satisfy (2.6) (and similarly (2.7)) almost surely for all $z \in \mathcal{C}$.

Since μ_z and $\tilde{\mu}_z$ are Stieltjes kernels, one can easily prove that $z \mapsto \int \frac{g}{D(\tilde{\mu}_z,\mu_z)} dH$ is analytic over $\mathbb{C}^+$. Therefore, by (2.6), the two analytic functions $z \mapsto \int g\, d\mu_z$ and $z \mapsto \int \frac{g}{D(\tilde{\mu}_z,\mu_z)} dH$ coincide almost surely over $\mathcal{C}$ which contains a limit point. They must be equal almost surely over $\mathbb{C}^+$. Therefore μ_z and $\tilde{\mu}_z$ satisfy (2.6) (and similarly (2.7)) almost surely for all $z \in \mathbb{C}^+$.

Since μ and $\tilde{\mu}$ are Stieltjes kernels satisfying almost surely (2.6) and (2.7), they must be almost surely equal to the unique pair of solutions $(\pi, \tilde{\pi})$ by Theorem 2.3. In particular, μ and $\tilde{\mu}$ are almost surely non-random. Thus for every subsequence $M = M(n)$,

$$\text{a.s.,} \quad \forall z \in \mathbb{C}^+, \quad L_z^M \xrightarrow[n\to\infty]{w} \pi_z \quad \text{and} \quad \tilde{L}_z^M \xrightarrow[n\to\infty]{w} \tilde{\pi}_z.$$

Therefore, the convergence remains true for the whole sequences L_z^n and $\tilde{L}_z^n$. Theorem 2.4 is proved.

5. Further Results and Remarks

In this section, we present two corollaries of Theorems 2.3 and 2.4. We will discuss the case where $\Lambda_n = 0$ and the case where the variance profile $\sigma(x, y)$ is constant. These results are already well-know ([4, 5, 8, 9]).

5.1. **The Centered case.**

Corollary 5.1. *Assume that (A-1) and (A-2) hold. Then the empirical distribution of the eigenvalues of the matrix* $Y_n\, Y_n^T$ *converges a.s. to a non-random probability measure* $\mathbb{P}$ *whose Stieltjes transform* f *is given by*

$$f(z) = \int_{[0,1]} \pi_z(dx),$$

where π_z *is the unique Stieltjes kernel with support included in* $[0, 1]$ *and satisfying*

$$\forall g \in C([0,1]), \qquad \int g\, d\pi_z = \int_0^1 \frac{g(u)}{-z + \int_0^1 \frac{\sigma^2(u,t)}{1+c\int_0^1 \sigma^2(x,t)\, \pi_z(dx)} dt} du. \tag{5.1}$$

Remark 5.1. In this case, one can prove that π_z is absolutely continuous with respect to du, i.e. $\pi_z(du) = k(z, u)du$ where $z \mapsto k(z, u)$ is analytic and $u \mapsto k(z, u)$ is continuous. Eq. (5.1) becomes

$$\forall u \in [0,1],\ \forall z \in \mathbb{C}^+, \quad k(u,z) = \frac{1}{-z + \int_0^1 \frac{\sigma^2(u,t)}{1+c\int_0^1 \sigma^2(x,t)\, k(x,z) dx} dt}. \tag{5.2}$$

Eq. (5.2) appears (up to notational differences) in [8] and in [4] in the setting of Gram matrices based on Gaussian fields.

Proof. Assumption (A-3) is satisfied with $\Lambda_{ii}^n = 0$ and $H(du, d\lambda) = du \otimes \delta_0(\lambda)$ where du denotes Lebesgue measure on $[0, 1]$. Therefore Theorems 2.3 and 2.4 yield the existence of kernels π_z and $\tilde{\pi}_z$ satisfying (2.6) and (2.7). It is straightforward to check that in this case π_z and $\tilde{\pi}_z$ do not depend on variable λ. Therefore (2.6) and (2.7) become:

$$\int g\, d\pi_z = \int \frac{g(u)}{-z(1 + \int \sigma^2(u,t)\tilde{\pi}(z,dt))} du \tag{5.3}$$

and

$$\begin{aligned} \int g\, d\tilde{\pi}_z \quad &= \quad c\int_{[0,1]} \frac{g(cu)}{-z(1 + c\int \sigma^2(t,cu)\pi(z,dt))} du \\ &\qquad +(1-c)\int_{[c,1]} \frac{g(u)}{-z(1 + c\int \sigma^2(t,u)\pi(z,dt))}\, du \\ &= \quad \int_{[0,1]} \frac{g(u)}{-z(1 + c\int \sigma^2(t,u)\pi(z,dt))} du, \end{aligned} \tag{5.4}$$

where $g \in C([0, 1])$. Replacing $\int \sigma^2(u, t)\tilde{\pi}(z, dt)$ in (5.3) by the expression given by (5.4), one gets the following equation satisfied by $\pi_z(du)$:

$$\int g\, d\pi_z = \int \frac{g(u)}{-z + \int \frac{\sigma^2(u,t)}{1+c\int \sigma^2(s,t)\pi(z,ds)} dt} du$$

□

5.2. **The non-centered case with i.i.d. entries.**

Corollary 5.2. *Assume that (A-1) and (A-2) hold where* $\sigma(x,y)=\sigma$ *is a constant function. Assume moreover that* $\frac{1}{N}\sum_{i=1}^{N}\delta_{\Lambda_{ii}^2}\to H_\Lambda(d\lambda)$ *weakly, where* H_Λ *has a compact support. Then the empirical distribution of the eigenvalues of the matrix* $\Sigma_n\Sigma_n^T$ *converges a.s. to a non-random probability measure* $\mathbb{P}$ *whose Stieltjes transform is given by*

$$f(z)=\int\frac{H_\Lambda(\,d\lambda)}{-z(1+c\sigma^2 f(z))+(1-c)\sigma^2+\frac{\lambda}{1+c\sigma^2 f(z)}}. \tag{5.5}$$

Remark 5.2. Eq. (5.5) appears in [5] in the case where $\Sigma_n=\sigma Z_n+R_n$ where Z_n and R_n are assumed to be independent, $Z_{ij}^n=\frac{X_{ij}}{\sqrt{n}}$, the X_{ij} being i.i.d. and the empirical distribution of the eigenvalues of $R_nR_n^T$ converging to a given probability distribution. Since R_n is not assumed to be diagonal in [5], the results in [5] do not follow from Corollary 5.2.

Proof. One can build a sequence $(i/n,\Lambda_{ii}^2)$ such that $\frac{1}{n}\sum_{i=1}^{n}\delta_{(i/n,\Lambda_{ii}^2)}\xrightarrow{\mathcal{D}} du\otimes H_\Lambda(d\lambda)$. Therefore (A-3) is satisfied with $H(du,d\lambda)=du\otimes H_\Lambda(d\lambda)$ and Theorems 2.3 and 2.4 yield the existence of kernels π_z and $\tilde\pi_z$ satisfying (2.6) and (2.7). It is straightforward to check that in this case π_z and $\tilde\pi_z$ do not depend on variable u. Equation (2.6) becomes

$$\int g\,d\pi_z=\int\frac{g(u,\lambda)}{-z(1+\sigma^2\int\tilde\pi(z,dt,d\zeta))+\frac{\lambda}{1+c\sigma^2\int\pi(z,dt,d\zeta)}}H(du,d\lambda)$$

Let $g(u,\lambda)=1$, then (2.6) becomes

$$f(z)=\int\frac{1}{-z(1+\sigma^2\tilde f(z))+\frac{\lambda}{1+c\sigma^2 f(z)}}H_\Lambda(d\lambda) \tag{5.6}$$

Denote by $f_n(z)=\frac{1}{N}\sum_1^N q_{ii}(z)$ and by $\tilde f_n(z)=\frac{1}{n}\sum_1^N\tilde q_{ii}(z)=\frac{1}{n}\mathrm{Tr}(\Sigma_n^T\Sigma_n-zI)^{-1}$. Since $f_n(z)=\frac{1}{N}\mathrm{Tr}(\Sigma_n\Sigma_n^T-zI)^{-1}$ and $\tilde f_n(z)=\frac{1}{n}\mathrm{Tr}(\Sigma_n^T\Sigma_n-zI)^{-1}$ (recall that $N\le n$), we have $\tilde f_n(z)=\frac{N}{n}f_n(z)+\left(1-\frac{N}{n}\right)\left(-\frac{1}{z}\right)$. This yields $\tilde f(z)=cf(z)-\frac{1-c}{z}$. Replacing $\tilde f(z)$ in (5.6) by this expression, we get (5.5). □

5.3. **Statement of the results in the complex case.**

In the complex setting, Assumptions (A-1)-(A-3) must be modified in the following way:

Assumption A-1. *The random variables* $(X_{ij}^n\ ;\ 1\le i\le N,\ 1\le j\le n\ ,\ n\ge 1)$ *are complex, independent and identically distributed. They are centered with* $\mathbb{E}|X_{ij}^n|^2=1$ *and satisfy:*

$$\exists\ \epsilon>0,\quad \mathbb{E}|X_{ij}^n|^{4+\epsilon}<\infty.$$

Assumption A-2. *The complex function* $\sigma:[0,1]\times[0,1]\to\mathbb{C}$ *is such that* $|\sigma|^2$ *is continuous and therefore there exist a non-negative constant* $\sigma_{\max}$ *such that*

$$\forall(x,y)\in[0,1]^2,\quad 0\le|\sigma(x,y)|^2\le\sigma_{\max}^2<\infty. \tag{5.7}$$

If Λ_n is a complex deterministic $N\times n$ matrix whose non-diagonal entries are zero, assume that:

Assumption A-3. *There exists a probability measure* $H(\,du,d\lambda)$ *over the set* $[0,1]\times\mathbb{R}$ *with compact support* $\mathcal{H}$ *such that*

$$\frac{1}{N}\sum_{i=1}^{N}\delta_{\left(\frac{i}{N},|\Lambda_{ii}^n|^2\right)}(du,d\lambda)\xrightarrow[n\to\infty]{\mathcal{D}}H(\,du,d\lambda). \tag{5.8}$$

In Theorem 2.3 (Eq. (2.6) and (2.7)), one must replace σ by its module $|\sigma|$. The statements of Theorem 2.4 and Corollary 2.5 are not modified.

References

[1] G. Anderson and O. Zeitouni. A CLT for a band matrix model. to be publihed in Probab. Theory Related Fields, 2005.

[2] Z. D. Bai. Methodologies in spectral analysis of large-dimensional random matrices, a review. *Statist. Sinica*, 9(3):611–677, 1999.

[3] Z. D. Bai and J. W. Silverstein. No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *Ann. Probab.*, 26(1):316–345, 1998.

[4] A. Boutet de Monvel, A. Khorunzhy, and V. Vasilchuk. Limiting eigenvalue distribution of random matrices with correlated entries. *Markov Process. Related Fields*, 2(4):607–636, 1996.

[5] R. Brent Dozier and Silverstein J. W. On the empirical distribution of eigenvalues of large dimensional information-plus-noise type matrices. preprint 2004.

[6] C.N. Chuah, J.M. Kahn, D.N.C. Tse, and R.A. Valenzuela. Capacity scaling in mimo wireless systems under correlated fading. *IEEE Trans. Inf. Theo.*, 48(3):637–650, 2002.

[7] M. Debbah, W. Hachem, P. Loubaton, and M. de Courville. MMSE analysis of certain large isometric random precoded systems. *IEEE Trans. Inform. Theory*, 49(5):1293–1311, 2003.

[8] V. L. Girko. *Theory of random determinants*, volume 45 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1990.

[9] V. L. Girko. *Theory of stochastic canonical equations. Vol. I*, volume 535 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 2001.

[10] A. Goldsmith, S.A. Jafar, N. Jindal, and S. Vishwanath. Capacity limits of mimo channels. *IEEE J. Sel. Areas in Comm.*, 21(5), 2003.

[11] T. Kailath, Sayed A.H., and Hassibi B. *Linear Estimation*. Prentice Hall, NJ, 2000.

[12] A. Khorunzhy, B. Khoruzhenko, and L. Pastur. Asymptotic properties of large random matrices with independent entries. *J. Math. Phys.*, 37(10):5033–5060, 1996.

[13] L. Li, A.M. Tulino, and S. Verdu. Design of reduced-rank mmse multiuser detectors using random matrix methods. *IEEE Trans. Inform. Theory*, 50(6):986–1008, june 2004.

[14] V. A. Marčenko and L. A. Pastur. Distribution of eigenvalues in certain sets of random matrices. *Mat. Sb. (N.S.)*, 72 (114):507–536, 1967.

[15] J. W. Silverstein. Strong convergence of the empirical distribution of eigenvalues of large-dimensional random matrices. *J. Multivariate Anal.*, 55(2):331–339, 1995.

[16] J. W. Silverstein and Z. D. Bai. On the empirical distribution of eigenvalues of a class of large-dimensional random matrices. *J. Multivariate Anal.*, 54(2):175–192, 1995.

[17] J. W. Silverstein and P.L. Combettes. Signal detection via spectral theory of large dimensional random matrices. *IEEE Trans. Signal Proces.*, 40(8):2100–2105, 1992.

[18] D. Tse and S. Hanly. Linear multiuser receivers: effective interference, effective bandwidth and user capacity. *IEEE Trans. Inform. Theory*, 45(2):641–657, 1999.

[19] D. Tse and O. Zeitouni. Linear multiuser receivers in random environments. *IEEE Trans. Inform. Theory*, 46(1):171–188, 2000.

[20] A. Tulino and S. Verdu. *Random matrix theory and wireless communications*, volume 1 of *Fondations and Trends in Communications and Information Theory*. Now Publishers, Delft, 2004.

[21] Y. Q. Yin. Limiting spectral distribution for a class of random matrices. *J. Multivariate Anal.*, 20(1):50–68, 1986.

Walid Hachem,
Supélec (Ecole Supérieure d'Electricité)
Plateau de Moulon, 3 rue Joliot-Curie
91192 Gif Sur Yvette Cedex, France.
e-mail: walid.hachem@supelec.fr

Philippe Loubaton,
IGM LabInfo, UMR 8049, Institut Gaspard Monge,
Université de Marne La Vallée, France.

5, Bd Descartes, Champs sur Marne,
77454 Marne La Vallée Cedex 2, France.
e-mail: loubaton@univ-mlv.fr

Jamal Najim,
CNRS, Télécom Paris
46, rue Barrault, 75013 Paris, France.
e-mail: najim@tsi.enst.fr